

Bergische Universität Wuppertal

Fakultät für Mathematik und Naturwissenschaften

Institute of Mathematical Modelling, Analysis and Computational Mathematics (IMACM)

Preprint BUW-IMACM 24/02

Manh Tuan Hoang and Matthias Ehrhardt

Differential equation models for infectious diseases: Mathematical modeling, qualitative analysis, numerical methods and applications

February 15, 2024

http://www.imacm.uni-wuppertal.de

Differential equation models for infectious diseases: Mathematical modeling, qualitative analysis, numerical methods and applications

Manh Tuan Hoang and Matthias Ehrhardt

ABSTRACT. Mathematical epidemiology has a long history of origin and development. In particular, mathematical modeling and analysis of infectious diseases has become a fundamental and indispensable approach to discovering the characteristics and mechanisms of the transmission dynamics of epidemics, thereby effectively predicting possible scenarios in reality, as well as controlling and preventing diseases.

In recent decades, differential equations have been widely used to model many important infectious diseases. The study of these differential equation models is very useful in both theory and practice, especially in proposing appropriate strategies for disease control and prevention. This is of great benefit to public health and health care.

In this survey article, we review many recent developments and real-life applications of deterministic differential equation models in modeling major infectious diseases, focusing on the following aspects: mathematical modeling, qualitative analysis, numerical methods, and real-life applications. We also present and discuss some open problems and future directions that research in differential equation models for infectious diseases can take.

This article is a good introduction to the topic, also to learn more about nonstandard finite difference (NSFD) methods.

1. Introduction

Infectious diseases have always been a major and constant threat to public health. Mankind has always had to face and fight many infectious diseases with varying degrees of danger, such as influenza, hepatitis, Zika, malaria, measles, tuberculosis, hepatitis, vector-borne diseases, Ebola, and most recently the COVID-19 pandemic. As a result, several approaches, policies, and strategies have been proposed to combat infectious diseases. Among these, mathematical epidemiology has a long history of origin and development. In particular, mathematical modeling and analysis of infectious diseases has become a fundamental and indispensable approach to discover the characteristics and transmission mechanisms of epidemics,

²⁰²⁰ Mathematics Subject Classification. Primary 34A34, 35Q92, 37N25, 37N30, 92B05, 65L05, 65L12, 92Dxx .

Key words and phrases. Mathematical modeling, Epidemiology, Epidemics, Infectious diseases, Numerical methods.

to effectively predict possible scenarios in reality, to control and prevent diseases [25, 70, 71, 72, 73, 74, 75, 297, 332].

The well-known SIR model, proposed by Kermack and McKendrick in 1927 [230], can be considered one of the first epidemic models and is usually used to introduce epidemic modeling. The study of mathematical models of infectious diseases is very useful in both theory and practice, especially in proposing appropriate strategies for disease control and prevention. This is of great benefit to public health and health care.

It is well known that differential equations, including ordinary differential equations (ODEs) and partial differential equations (PDEs), have several useful applications in real life. They are widely used to describe many important phenomena and processes in science and engineering (see e.g. [25, 49, 70, 71, 72, 73, 74, 75, 234, 297, 332, 413]). One of its prominent applications is the mathematical modeling and analysis of infectious diseases. Over the past few decades, a large number of differential equation models have been extensively developed to explore the transmission dynamics of major infectious diseases. These models have confirmed the important role of differential equations in epidemic modeling.

Nowadays, epidemic models based on differential equations have always been an important and indispensable approach in modeling infectious diseases, especially in the context that epidemics are constantly changing and posing new challenges. For differential equation models of infectious diseases, the following aspects are mainly focused:

- *Mathematical Modeling:* The use of differential equations and the foundations of mathematical epidemiology to propose mathematical models that describe the transmission of infectious diseases.
- *Qualitative study:* Investigate mathematical properties of the proposed differential equation models, including existence and uniqueness of solutions, positivity and boundedness of solutions, asymptotic stability properties, conservation laws, physical properties, and basic reproduction number.
- *Numerical Methods:* Construction of efficient numerical methods, especially numerical methods that preserve important mathematical features of the proposed differential equation models.
- *Practical Applications:* Applying the theoretical results to provide scenarios of disease spread, to suggest anti-epidemic measures and strategies, to evaluate the effectiveness of vaccines and existing anti-epidemic measures, to study the spread of computer viruses, rumors and malware on the Internet, and to model animal diseases. and animal disease modeling with applications in agriculture.

The aim of this review article is to review many recent developments and reallife applications of deterministic differential equation models in modeling major infectious diseases, focusing mainly on the following aspects: mathematical modeling, qualitative analysis, numerical methods, and real-life applications. We also present and discuss some open problems and future directions that research in differential equation models for infectious diseases can take.

The outline of this article is as follows: In Section 2, we provide an overview of epidemic models based on differential equations, considering basic models and their variants and extensions. In Section 3 we focus on the qualitative analysis aspect and

its practical applications. Numerical methods are presented in Section 4. Future research and open problems are discussed in Section 5. The last section contains concluding remarks and discussions.

2. Mathematical Modeling

In this section, we review results on mathematical modeling based on deterministic ODEs and PDEs for infectious diseases.

2.1. ODE models: Basic Epidemic Models. We start with one of the first and basic epidemic models introduced by Kermack and McKendrick in 1927 [230]. For this purpose, let us consider general autonomous dynamical systems described by ODEs of the form

(2.1)
$$\dot{y}(t) = f(y(t)), \quad t > 0, \qquad y(0) = y_0 \in \mathbb{R}^n,$$

where $y = [y_1, y_2, \ldots, y_n]^\top : [0, \infty) \to \mathbb{R}^n$, $f = [f_1, f_2, \ldots, f_n]^\top : \mathbb{R}^n \to \mathbb{R}^n$ and \dot{y} stands for the time derivative of y. Here it is assumed that the right-hand-side function f satisfies all necessary smoothness assumptions so that solutions of (2.1) exist and are unique (see e.g. [49, 234, 413]).

Many mathematical models based on (2.1) have been proposed to study epidemic models. In these models, diseases caused by viruses or bacteria are not modelled directly in the population model, but only indirectly through the number of infected individuals. For example, the classical SI, SIS and SIR epidemic models classify individuals in the population according to their status with respect to the disease: healthy, infected and immune. More clearly, the disease states S, I and R are defined as follows [25, 439]:

- *susceptible S*: Individuals who are not infected but are susceptible to acquiring the disease and becoming contagious.
- *infected I*: Individuals who have been infected, are currently contagious, and have the potential to spread the disease to others.
- *removed R*: Individuals who have experienced the disease, recovered, and achieved permanent immunity, or are isolated until both recovery and permanent immunity are achieved.

Models with these states are called *SIR models*, adapted to the characteristics of the infectious disease, for example:

- SI implies the absence of any possible recovery: $S \rightarrow I$;
- SIS indicates the possibility of recovery, but does not guarantee immunity: $S \rightarrow I \rightarrow S$;
- SIR represents a temporary state of immunity: $S \to I \to R \to S$.

One of the simplest models involves the dynamics of S-, I-, R- individuals, first introduced by Kermack and McKendrick in 1927 [230] (see also [297]):

(2.2)

$$\dot{S}(t) = -\beta I(t)S(t),$$

$$\dot{I}(t) = \beta I(t)S(t) - \alpha I,$$

$$\dot{R}(t) = \alpha I(t), \quad t > 0,$$

where

- β is the proportionality constant ('transmission rate');
- α is the recovery rate;

- $\beta I(t)$ is called the force of infection.
- βSI represents the number of new infections per unit of time (incidence).

Although the SIR model (2.2) looks analytically simple, finding its exact analytical solution is an interesting problem. Some analytical techniques used to find the solution of (2.2) can be found in [92, 189, 237].

It is not difficult to analyze basic mathematical properties of the Kermack-McKendrick SIR model [230, 297]. More clearly, it can be shown that

$$\lim_{t \to \infty} S(t) = S_{\infty} > 0, \quad \lim_{t \to \infty} R(t) = R_{\infty} > 0, \quad \lim_{t \to \infty} I(t) = I_{\infty} = 0.$$

The quantity S_{∞} is called the *final size of the epidemic*. In particular, the function I(t) of infected individuals can monotonically decrease to zero, or first monotonically increase to some maximum value I_{\max} and then decrease to zero. Here, a necessary and sufficient condition for the initial increase of I(t) is easily determined and is given by

$$S(0) > \frac{\alpha}{\beta}.$$

On the other hand, I_{max} can be computed as

$$I_{\max} = -\frac{\alpha}{\beta} + \frac{\alpha}{\beta} \ln \frac{\alpha}{\beta} + S_0 + I_0 - \frac{\alpha}{\beta} \ln S_0.$$

The quantity I_{max} is very useful in estimating the progression of epidemics since it indicates when the number of infections will begin to decline.

Note that the Kermack-McKendrick SIR epidemic model, for example, uses some hypotheses:

- Infected individuals are also infectious;
- the total population remains constant;
- the population experiences no births or deaths;
- the population is closed, that is, no outside individuals enter or leave the population;
- all recovered individuals have complete immunity and are impervious to reinfection.

The above assumptions may seem rather restrictive, but they can be satisfied within certain limits. For example, several childhood diseases such as chickenpox, smallpox, rubella, mumps, scarlet fever, hand-foot-and-mouth disease lead to permanent immunity, or many vaccines can create long-lasting or even lifelong immunity [297].

Although the Kermack-McKendrick SIR epidemic model is simple and under some strict assumptions, it is still appropriate and effective for modeling many infectious diseases. In fact, once we have given disease-specific time series data, the parameter estimation problem for the SIR model can be solved by comparing its solution to the given data. Examples of parameter estimation from data can be found in [70, 72, 297]. Recently, the Kermack-McKendrick SIR epidemic model was used to study and predict the transmission dynamics of the COVID-19 pandemic [228, 244, 254, 296, 331, 401, 438].

In [231], the limitation of the SIR model (2.2) was improved by considering the effect of the continuous introduction of new susceptible individuals into the population. However, the results presented in [231] had two important limitations. One was that the disease of interest was the only cause of death, and the second was that the age of the individuals did not affect their infectivity, susceptibility, or reproductive capacity. In [232], the first of the above limitations was overcome

by the introduction of constant non-specific mortality rates, which, for the sake of generality, are assumed to be different. are assumed to be different for virgins (individuals who have never been infected), sick, and recovered.

In general, the classical SIR model should be adapted to the characteristics of each epidemic.

2.2. Variants and Extensions of the Basic Models. The classical epidemic models have played an important role in epidemic modeling. Inspired by basic epidemic models and principles of mathematical epidemiology, many mathematical models models have been proposed and developed to study infectious diseases.

There are several types of incidence, depending on the assumption made about the force of infection. One of the simplest forms is the mass action incidence or bilinear incidence function, which is $f(S, I) = \beta SI$. In the model (2.2), the *interaction term* βIS is a linearly increasing function of the number of infected individuals. As analyzed in [90], while this interaction term may be true for small I, it seems rather unrealistic that it can still hold for large I. For this reason, Capasso and Serio modified (2.2) by replacing the linear interaction term βIS by a non-linear function g(I)S, where g(I) satisfies

- (1) $\forall x \in \mathbb{R}_+ : g(x) \ge 0;$
- (2) g(0) = 0;
- (3) $\exists c \in \mathbb{R}_+ \setminus \{0\}$ s.t. $\forall x \in \mathbb{R}_+ : g(x) \le c;$
- (4) $g'(x) \colon \mathbb{R}_+ \to \mathbb{R}$, the derivative of g, exists and is bounded on any compact interval of \mathbb{R}_+ , with g'(0) > 0;
- (5) $\forall x \in \mathbb{R}_+ : g(x) \le xg'(0)$, where $\mathbb{R}_+ := [0, \infty)$.

The function g(I) takes into account the "saturation" phenomenon or the other "psychological" effects. Two famous nonlinear incidence functions are the saturated incidence rate $f(S, I) = \beta SI/(1 + \gamma I)$ and the standard incidence function $f(S, I) = \beta SI/(S+I)$. Epidemic models using generalized nonlinear incidence rate can be found in [151, 152, 165, 192, 212, 261, 270, 272, 303, 378, 399, 424].

In the SIR model, it was assumed (2.2) that the rate of contacts per infective is proportional to the total population size N, which was widely used in all early epidemic models. As mentioned in [70, 74], this assumption is quite unrealistic except in the early stages of an epidemic occurring within a moderately sized population. It is more realistic to consider a contact rate that is a non-increasing function of total population size. The SIR model can then be generalized by assuming that an average member of the population makes C(N) contacts per unit time, with $C'(N) \geq 0$, and defining

$$\beta(N) = \frac{C(N)}{N},$$

where $\beta'(N)$ is assumed to be negative to express the idea of saturation in the number of contacts. The following are some special cases of C(N) that have been widely used in epidemic modeling with general contact rates.

- Standard incidence: $C(N) = \lambda;$
- Mass action incidence: $C(N) = \beta N$;
- Interaction of Michaelis-Menten type:

$$C(N) = \frac{aN}{1+bN},$$

which was used in [134].

• Saturating contact rate based on a mechanistic derivation for pair formation [190]

$$C(N) = \frac{aN}{1 + bN + \sqrt{1 + 2bN}}$$

• $C(N) = \lambda N^{\alpha}$ with $\alpha = 0.05$ was used in [306]. It has been shown that this function works quite well for data on contact-borne diseases in medium-sized cities.

In recent decades, the basic classical epidemic models and their variants have been extensively developed to describe the transmission dynamics of many major infectious diseases:

- Basic virus dynamics models [30, 67, 149, 196, 340, 427];
- Influenza [6, 8, 93, 182, 394];
- Severe acute respiratory syndrome (SARS) [69, 110, 179, 210, 337, 469];
- Ebola [57, 130, 264, 291, 335];
- Hepatitis B and C [154, 197, 202, 293, 298, 324, 339, 358, 411, 416, 446, 459, 466];
- Tuberculosis [64, 181, 275, 333, 372, 405, 422, 425];
- Vector-borne diseases [62, 77, 117, 217, 263, 381, 450];
- Malaria [14, 161, 216, 240, 289, 338, 386, 420, 428];
- Measles [16]
- Zika virus [9, 163, 239, 285, 336, 392, 448, 470];
- Dengue fever [12, 97, 142, 336, 360, 366];
- COVID-19 pandemic [10, 15, 28, 58, 118, 141, 203, 228, 238, 241, 244, 254, 259, 288, 296, 327, 330, 331, 345, 353, 370, 375, 401, 420, 438];
- HIV/AIDS [155, 178, 221, 273, 280, 341, 379, 434].

Besides, epidemic models are widely used in

- Diabetes Mellitus [95, 333];
- cancer: malignant invasion of tumor cells [295];
- cervical cancer: human papillomavirus model [81];
- animal disease modeling with applications in agriculture [1, 7, 44, 427];
- chemostat models to represent microbial growth and competition [17, 18, 403];
- modeling the spreading of computer viruses and rumors on the Internet [165, 211, 219, 268, 355, 356, 367, 368, 390, 462, 463, 464, 474, 475];
- modeling addictions, e.g. alcohol drinking [121, 214, 382, 456], tobacco [167, 260, 274, 393, 433, 440], heroin [113, 274, 328, 410, 451], opioids [55, 80, 87, 113, 472], cocaine [383, 385], drug consumption [129, 173, 441], obesity [45, 85, 150, 223, 384], etc.

It should be emphasized that the ODE models of the form (2.1) are also extended in the context of delayed systems [5, 94, 115, 149, 159, 160, 195, 350, 371, 404, 429, 458]; time fractional-order systems [5, 32, 105, 414, 427]; and stochastic systems [24, 27, 31, 76, 84, 104, 174, 389] for modeling infectious diseases. These extended models provide an additional powerful approach to the disease analysis.

2.3. PDE Models. In addition to ODE models of the form (2.1), PDE models, which extend ODE models, have also been extensively studied for the analysis of infectious diseases [25, 71, 75, 297, 332, 387].

More specifically, compartmental models in epidemiology can be extended by using spatial reaction-diffusion systems, where each compartment, representing a different species, is allowed to invade a spatial domain $\Omega \subset \mathbb{R}^m$ (or a metric graph network) with a space-dependent density. The densities interact with each other according to the same mathematical laws as for the space-independent case, but are individually subject to a spatial diffusion mechanism, usually associated with the Laplace operator [42]. Then a system of n interacting species, each with a spatial density

$$\{u_i(x,t): x \in \Omega, \quad t \ge 0\}, \quad i = 1, 2, \dots, n$$

can be described by a system of semilinear parabolic PDEs of the form

(2.3)
$$\frac{\partial u}{\partial t}(x,t) = D\Delta u(x,t) + f(u(x,t))$$

supplied with suitable boundary conditions, where $D = \text{diag}(d_1, d_2, \dots, d_n), f \colon \mathbb{R}^n \to \mathbb{R}$ is the interaction law among the species via their densities, and

$$\Delta u(x,t) = \frac{\partial^2 u}{\partial x_1^2}(x,t) + \ldots + \frac{\partial^2 u}{\partial x_n^2}(x,t).$$

Spatial models of the form (2.3) have been used to study the transmission of infection, depending on how a particular disease is transmitted between different populations or subpopulations.

Allen et al. $[{\bf 26}]$ proposed an SIS reaction-diffusion model in a heterogeneous environment in the form

(2.4)
$$\begin{aligned} &\frac{\partial}{\partial t}S(t,x) = d_S \Delta S(t,x) - \frac{\beta(x)S(t,x)I(t,x)}{S(t,x) + I(t,x)} + \gamma(x)I(t,x), \quad t > 0, \ x \in \Omega, \\ &\frac{\partial}{\partial t}I(t,x) = d_I \Delta I(t,x) + \frac{\beta(x)S(t,x)I(t,x)}{S(t,x) - I(t,x)} - \gamma(x)I(t,x), \quad t > 0, \ x \in \Omega, \end{aligned}$$

with the coupling condition

(2.5)
$$\frac{\partial}{\partial \mathbf{n}} S(t, x) = \frac{\partial}{\partial \mathbf{n}} I(t, x) = 0,$$

where

- S(t, x) and I(t, x) denote the density of susceptible and infectious individuals in a given spatial region Ω , which is assumed to be a bounded domain in $\mathbb{R}^n (n \ge 1)$ with a smooth boundary $\partial \Omega$;
- Ω is isolated from the outside for the host, implying the homogeneous Neumann boundary condition; **n** is the outward unit normal vector on $\partial\Omega$, and $\partial/\partial \mathbf{n}$ denotes the normal derivative along **n** on $\partial\Omega$.
- d_S and d_I are the dispersion for susceptible and infectious individuals, respectively;
- the positive functions $\beta(x)$ and $\gamma(x)$ are the spatially dependent transmission and recovery rates at position $x \in \Omega$, respectively.

The existence, uniqueness and asymptotic profile of the equilibria are then analyzed.

In [213], Huang et al. proposed and studied two modified SIS diffusion models associated with the Dirichlet boundary condition, which reflects a hostile environment in the boundary. The analysis of the basic reproduction number and a partial result on the global stability of the endemic equilibrium are presented.

In [255], a spatially diffusive SIR epidemic model with the mass action infection mechanism and homogeneous Neumann boundary condition was considered in the form

$$\begin{aligned} &(2.6)\\ &\frac{\partial}{\partial t}S(t,x) = k_S \Delta S(t,x) + b(x) - \beta(x)S(t,x)I(t,x) - \mu(x)S(t,x), \quad t > 0, \ x \in \Omega, \\ &\frac{\partial}{\partial t}I(t,x) = k_I \Delta I(t,x) + \beta(x)S(t,x)I(t,x) + (\mu(x) + \gamma(x))I(t,x), \quad t > 0, \ x \in \Omega, \\ &\frac{\partial}{\partial t}R(t,x) = k_R \Delta R(t,x) + \gamma(x)I(t,x) - \mu(x)R(t,x), \quad t > 0, \quad x \in \Omega, \end{aligned}$$

with initial data

(2.7)
$$S(0,x) = S_0(x), \quad I(0,x) = I_0(x), \quad R(0,x) = R_0(x), \quad x \in \Omega,$$

and coupling conditions

(2.8)
$$\frac{\partial}{\partial \mathbf{n}} S(t,x) = \frac{\partial}{\partial \mathbf{n}} I(t,x) = \frac{\partial}{\partial \mathbf{n}} R(t,x)0,$$

where

- S(t, x), I(t, x) and R(t, x) denote the populations of susceptible, infective and recovered individuals at position x and time t, respectively;
- k_S , k_I and k_R denote the dissemination rates for susceptible, infectious and recovered individuals, respectively;
- b(x), β(x), μ(x) and γ(x) denote the birth rate, the transmission rate, the mortality rate and the recovery rate at position x, respectively.

By discretizing the PDE model (2.6) with respect to the space variable and constructing Lyapunov functions for the corresponding ODE models, the global asymptotic stability of (2.6) has been established [255].

In [256], the model (2.6) is extended by a new more realistic model with non-local diffusion.

In a recent paper, some extensions of the classical SIR model with non-symmetric spatial dependence are introduced to study the spread of some diseases [419].

Also, a large number of spatial reaction-diffusion models of major infectious diseases such as HBV, malaria, influenza, West Nile virus transmission, Zika, etc. can be found in [50, 91, 108, 122, 233, 242, 243, 266, 276, 277, 326, 369, 396, 406, 415, 437, 443, 444, 445, 455, 457, 459, 473], in which the models proposed in [108, 233, 242, 276, 396, 437, 455, 473] can be directly used to study the COVID-19 epidemic.

3. Qualitative analysis and applications

Qualitative analysis of differential equations modeling infectious diseases is very important since it can have many useful applications in reality, such as suggesting appropriate strategies for disease control and prevention; evaluating the effects of vaccines; waning immunity; parameter estimation problems; parameter sensitivity analysis and optimal control strategies (usually w.r.t. vaccination strategies, stakeholder decisions (wearing masks, physical isolation, curfews, etc.).

In this section, we emphasize qualitative analytical aspects of differential equation models and their applications. Methods, approaches and tools used in qualitative analysis are also discussed.

3.1. Analysis of ODE models. The first property of interest for ODE models of infectious diseases is well-posedness, including existence, uniqueness of solutions, and continuous dependence on initial data. Well-posedness is easy to establish and is often automatically satisfied due to the smoothness of the right-hand-side functions [25, 49, 234, 413]. In general, in addition to well-posedness, qualitative analysis aspects of ODE models of infectious diseases focus mainly on the following issues.

3.1.1. The positivity and boundedness of the solutions. Obviously, positivity should be an obvious property of the solutions of ODE models for infectious diseases, i.e. $y(t) \in \mathbb{R}^n_+ = \{(y_1, y_2, \ldots, y_n) \in \mathbb{R}^n | y_1, y_2, \ldots, y_n \ge 0\}$ for t > 0 whenever $y(0) \in \mathbb{R}^n_+$. In this case, the set \mathbb{R}^n_+ is called a *positively invariant set*. This property can be easily verified using well-known theorems on the positivity of ODEs [**209**, Lemma 1], [**403**]. Meanwhile, boundedness can be established on the basis of comparison theorems for differential equations [**304**]. Note that positively invariant sets and feasible sets of ODE models are obtained from their positivity and boundedness.

3.1.2. Conservation laws. Many ODE models in population dynamics and also in epidemiology can satisfy some *conservation laws*, such as direct, generalized and subconservation laws [**317**, **320**]. Conservation laws for ODE models of infectious diseases can be established based on the theory of ODEs [**17**, **25**, **234**, **403**, **413**] or comparison theorems for differential equations [**304**].

3.1.3. Equilibrium points. Equilibrium points of ODE models of the form (2.1) are solutions of the equation f(y) = 0. An equilibrium point is also called a fixed point, constant solution, steady state, critical point or a steady-state solution [25, 234, 413]. In general, it is not difficult to determine the set of equilibrium points, except when the ODE model under consideration has high dimensions and contains many parameters. Two common types of equilibria are disease-free equilibrium (DFE) and endemic equilibrium (EE) points, which correspond to the possibility of the epidemic being suppressed or remaining in the community.

3.1.4. Local asymptotic stability (LAS). The LAS of equilibrium points can be studied by the Lyapunov direct method using the Routh-Hurwitz criteria [25, 234, 413]. This approach analyzes the LAS of an equilibrium point by considering the position of the eigenvalues of the Jacobian matrix evaluated at the equilibrium point with respect to the left-half plane. More specifically, an equilibrium point y^* is locally asymptotically stable if all eigenvalues λ of the Jacobian $J(y^*) =$ $(\partial f/\partial y)(y^*)$ satisfy $Re(\lambda) < 0$, and it is unstable if $Re(\lambda) > 0$ for one or more of the eigenvalues of J. Note that the direct Lyapunov method is only applicable to hyperbolic equilibrium points. Here, an equilibrium point y^* is said to be hyperbolic if none of the eigenvalues of the matrix J lie on the imaginary axis, and nonhyperbolic otherwise, cf. [413].

3.1.5. Global asymptotic stability (GAS). The GAS analysis of equilibrium points is a very important problem because it can reveal the future evolution of epidemics. In particular, the GAS of free-disease equilibrium points indicates that epidemics will be extinguished, while the GAS of endemic-equilibrium points indicates that epidemics will exist stably in the population. In general, the GAS problem is not an easy one. One of the most successful approaches to this problem is the Lyapunov stability theory [262, 284]. In particular, several classes of Lyapunov functions have been proposed to analyze the GAS of ODE models in epidemiology [89, 248, 249, 250, 344, 370, 397, 435, 460], where common classes of Lyapunov functions are linear, quadratic and Volterra-type Lyapunov functions for a given ODE model.

On the other hand, the geometric method is a remarkable approach to the GAS analysis of ODEs [269, 270, 271]. Also, the Poincare-Bendixson theorem in combination with the Bendixson-Dulac criterion is very useful in studying the GAS of two-dimensional dynamical systems governed by ODEs [25, 297].

3.1.6. Basic reproduction number. One of the most important concerns about any infectious disease is its reproductive number \mathcal{R}_0 , which is useful in guiding control strategies [133, 430, 431, 432]. The basic reproduction number can be defined as the expected number of secondary cases produced by a typical infected individual during its entire period of infectiousness in a fully susceptible population [133]. It can also be considered as a threshold parameter for the local asymptotic stability of the disease-free equilibrium [430]. The basic reproduction number of epidemic models with the help of sensitivity analysis is very useful in guiding control strategies.

3.1.7. Optimal control problems. Epidemic models based on differential equations are often combined with optimal control strategies to find effective disease control measures [52, 60, 78, 103, 236, 282, 392, 395]. Here, the proposed optimal control problems are solved using Pontryagin's maximum principle [357].

3.1.8. Epidemic models with effect of vaccines. It is well known that vaccines are effective tools to combat infectious diseases and to project people against diseases. For this reason, epidemic models with the effect of vaccines are often considered [6, 20, 143, 144, 149, 153, 166, 177, 180, 215, 253, 377, 398]. The study of vaccination models can evaluate the efficacy of certain vaccines and suggest effective vaccination strategies.

3.1.9. Parameter estimation problem. ODE models for infectious diseases can be combined with real data of diseases to predict possible scenarios in reality. Therefore, the parameter estimation problem is very important to find best-fit parameters [70, 72, 297]. Following this approach, the parameter estimation problem has been extensively studied for several epidemic models [111, 286, 329, 362, 380], especially for the COVID-19 pandemic [228, 244, 282, 296, 331].

3.1.10. Bifurcation analysis and chaos. It is well-known that bifurcation theory studies qualitative changes in the state of a system as a parameter is varied [96, 258]. In general, applications of bifurcation analysis in epidemiology are very diverse, especially in studying the evolution and determining factors that may be associated with the suppression or outbreak of disease. For example, the forward bifurcation phenomenon, first noted by Kermack and McKendrick in [230], can be observed in several disease transmission models [146]. For epidemic models that exhibit forward bifurcation, the condition $\mathcal{R}_0 < 1$ is a necessary and sufficient condition for disease elimination [146, 183]. For many years, bifurcation analysis

for epidemic models has been studied extensively with many useful applications, including forward bifurcation, *backward bifurcation*, *Hopf bifurcation*, *Bogdanov-Takens bifurcation*, *saddle-node bifurcation*, *flip bifurcation* are mainly focused [21, 23, 48, 68, 109, 170, 185, 222, 257, 279, 301, 376, 391].

Chaos theory has many useful applications in many fields such as physics, biology, ecology and epidemiology, economics, etc. [65, 198, 302, 388]. Over the past few decades, chaos theory has been developed and studied with the aim of discovering chaotic phenomena/dynamics, complicated or even unpredictable dynamical behavior in epidemic models [59, 66, 79, 147, 170, 175, 225, 290, 291, 292, 342, 343].

3.2. Analysis of PDE models. In general, the qualitative analysis aspects for PDE models of infectious diseases are very similar to those of ODE models. In particular, the qualitative analysis of PDE models also focuses on well-posedness of mathematical models, positivity and boundedness of the solution, conservation laws, equilibria and their asymptotic stability, basic reproduction numbers and their implications, optimal control problems, parameter estimation, vaccination models, bifurcations and chaos [26, 50, 53, 54, 83, 98, 99, 116, 117, 122, 145, 164, 172, 213, 220, 233, 243, 255, 256, 266, 267, 277, 281, 287, 325, 326, 352, 369, 396, 406, 407, 409, 415, 419, 437, 442, 443, 444, 445, 449, 455, 457, 459, 461, 468, 471, 473, 476]. Some methods and tools used in the qualitative analysis of ODE models, such as basic reproduction number, Lyapunov stability theory, optimal control, bifurcation and chaos analysis, can be developed and extended for PDE models. However, the qualitative study for PDE models is more challenging due to the complexity of their structures.

4. Numerical methods

4.1. Standard and nonstandard numerical methods. It is well known that both ODEs and PDEs can be solved exactly only in a small number of cases, and that in most real-world situations it is almost inevitable to find approximate solutions. For this reason, numerical methods for differential equations have become one of the most fundamental and practically important research tasks [49, 186, 187, 265, 402, 412, 413, 423].

Numerical solutions for ODE models can be easily obtained using standard numerical methods such as the Runge-Kutta and Taylor (one-step) methods and multistep methods, while finite difference methods are appropriate and efficient for the numerical solution of PDE models [49, 186, 187, 265, 404, 412, 413, 423]. However, mathematical models arising in real-world applications in general, and in infectious disease modeling in particular, often possess several essential qualitative features, such as positivity, boundedness, asymptotic stability properties, conservation laws, periodicity and physical properties, etc., which must be respected by corresponding numerical schemes. Therefore, an important requirement for numerical methods is that they correctly preserve the essential properties of the corresponding differential equations. However, it has been shown by Mickens in [308, 311, 315, 316, 319, 321] that standard numerical methods cannot preserve the mathematical properties of ODEs for all values of the temporal step size.

In the 1980s, Mickens proposed the concept of *nonstandard finite difference* (*NSFD*) methods to compensate for drawbacks and shortcomings of standard numerical methods [**308**, **311**, **315**, **316**, **319**, **321**]. One of the main and outstanding

advantages of NSFD methods is that they can preserve essential mathematical properties of differential equations independently of the values of the step size. Such NSFD methods are said to be *dynamically consistent*. Thus, dynamically consistent NSFD methods are efficient and suitable for simulating the behavior of dynamic differential equation models over long periods of time.

In addition to NSFD methods for ODEs, geometric numerical integration [88, 176, 188] (or both [85]) and positivity-preserving Runge-Kutta methods [61, 169, 209] and modified Patankar-Runge-Kutta schemes [246, 247] have also been developed to construct reliable numerical methods that preserve the positivity as well as other dynamical properties of ODE models.

In the next subsection, we provide an overview of NSFD methods for mathematical models of infectious diseases and their applications.

4.2. Nonstandard finite difference methods for epidemiological models of infectious diseases. In numerical analysis, *numerical instabilities* are solutions of finite difference models that do not correspond to any solution of the counterpart differential equation [**319**]. Mickens, the creator of the concept of NSFD methods, wrote: "Numerical instabilities are an indication that the discrete models are unable to model the correct mathematical properties of the solutions to the differential equations of interest" [**308**, **311**, **315**, **316**, **319**, **321**]. The concept of NSFD schemes was first introduced by Mickens in the 1980s to overcome the usual numerical instabilities associated with standard finite-difference schemes [**308**, **311**, **315**, **316**, **319**, **321**]. A finite difference scheme is said to be *nonstandard* if it is constructed based on a set of basic rules proposed by Mickens [**308**, **311**, **315**, **316**, **319**, **321**]. In particular, NSFD schemes for the ODE models of the form (2.1) can be defined as follows.

Consider a general finite difference scheme for (2.1) of the form

$$(4.1) D_{\Delta t}(y_k) = F_{\Delta t}(f; y_k),$$

where $D_{\Delta t}(y_k) \approx dy/dt$, $F_{\Delta t}(f; y_k) \approx f(y)$ and $t_k = k\Delta t$, Δt is the step size.

DEFINITION 4.1 ([**33**, **38**, **138**]). The finite difference scheme (4.1) is called an NSFD scheme if at least one of the following conditions is satisfied:

- $D_{\Delta t}(y_k) = \frac{y_{k+1} y_k}{\phi(\Delta t)}$, where $\phi(\Delta t) = \Delta t + \mathcal{O}(\Delta t^2)$ is a non-negative function and is called a nonstandard denominator function;
- $F_{\Delta t}(f; y_k) = g(y_k, y_{k+1}, \Delta t)$, where $g(y_k, y_{k+1}, \Delta t)$ is a non-local approximation of the right-hand side of the system (2.1).

NSFD schemes for (parabolic) PDEs [29, 101, 107, 132, 193, 227, 252, 305, 313, 334, 348, 359], fractional-order differential equations [82], delay differential equations are similarly defined based on the Mickens' methodology.

The main advantage of NSFD schemes over standard schemes is expressed in the following definitions.

DEFINITION 4.2 ([**33**, **38**]). Assume that the solutions of the equation (2.1) satisfy some property \mathcal{P} . The numerical scheme (4.1) is said to be (qualitatively) stable with respect to the property \mathcal{P} (or \mathcal{P} -stable), if for every value of $\Delta t > 0$ the set of solutions of (4.1) satisfies the property \mathcal{P} .

DEFINITION 4.3 ([35, 278, 315]). Consider the differential equation dy/dt = f(y). Let a finite difference scheme for the equation be $y_{k+1} = F(y_k; \Delta t)$. Let the

differential equation and/or its solutions have the property \mathcal{P} . The discrete model equation is dynamically consistent with the differential equation if it and/or its solutions also have the property \mathcal{P} .

Nowadays, NSFD methods based on the Mickens' methodology have become an efficient approach for numerically solving ODE models arising in real-world problems [3, 33, 34, 38, 119, 120, 123, 124, 125, 126, 127, 128, 135, 136, 137, 138, 139, 140, 158, 206, 207, 307, 308, 311, 312, 315, 316, 317, 319, 320, 321, 323, 349, 351, 373, 374, 408, 452, 453, 454]. In particular, NSFD schemes have been extensively studied for epidemic models, such as

- General epidemiological models [38, 46, 47, 100, 184, 299, 300]
- Influenza disease [162, 224, 235];
- Ebola [15, 39, 57, 218, 417];
- Hepatitis B [201, 202];
- Visceral Leishmaniasis [2, 400];
- Malaria [37, 156];
- Measles [11, 157];
- Zika [285, 421];
- COVID-19 [58, 118, 191, 194, 203, 288, 361, 426];
- Cancer: malignant invasion of tumor cells [43];
- Computer virus propagation models [127, 199, 367].

Compared to numerical methods for ODE models, numerical methods for PDE models are more challenging. Finite difference methods are one of the most common and efficient approaches for numerical simulation of PDEs [49, 265, 412, 423]. It is important to note that positivity should be an obvious property of the solutions of both ODE and PDE models for infectious diseases. Therefore, positivity preserving numerical methods are essential. To the best of our knowledge, numerical methods that preserve positivity and other dynamical properties for the PDE models are few. However, NSFD methods based on Mickens' methodology have been shown to be suitable and effective in constructing such numerical methods [33, 35, 36, 102, 114, 148, 308, 309, 310, 311, 314, 315, 316, 318, 319, 321, 322, 447]. In particular, dynamically consistent NSFD schemes have been applied to solve some PDE models of infectious diseases [168, 293, 294, 347, 358, 415, 416, 465].

Even though NSFD methods have several advantages, most of the existing dynamically consistent NSFD methods are only first-order convergent [106, 112, 119, 199, 201, 202], which can be considered as an inherent drawback of NSFD methods. For this reason, the problem of improving the accuracy of NSFD methods has attracted the attention of many researchers [19, 106, 126, 171, 200, 204, 205, 207, 208, 245, 299, 300]. However, it is very challenging to construct dynamically consistent NSFD methods, especially high-order methods, for differential equations.

In recent years, there has been an increased interest in solving PDEs using *Deep Learning* (see e.g. [56, 63, 191, 363]). More recently, in [251], a deep learning approach has been proposed to improve numerical methods for PDEs. This approach is based on an approximation of the local truncation error of the numerical method used to approximate the spatial derivatives of a given PDE.

In general, the construction of numerical methods, especially those that preserve important properties of differential models, is an important problem but not easy to solve. In addition, high-order numerical methods are still an important problem that has not been fully solved, and the reduced spatial accuracy of NSFD methods for PDEs is still an open problem.

NSFD methods for PDEs have lacked guaranteed first-order temporal accuracy and consistency for key models such as diffusion and reaction-diffusion systems. In a recent paper Pasha, Nawaz and Arif [347] proposed a novel NSFD scheme that overcomes this limitation and guarantees first-order temporal accuracy and second-order spatial accuracy while preserving positivity.

5. Future research and open problems

Although research on differential equation models for infectious diseases has been extensively developed over the past decades and has achieved many important successes, these models still need to be studied and expanded for the following reasons.

First, mankind is always facing and fighting many infectious diseases, which are not only constantly changing but also difficult to predict, and thus always pose a great and constant threat to public health. In this context, the development of mathematical models of infectious diseases remains a fundamental and effective approach to discover the characteristics and mechanisms of transmission of epidemics, and thus effectively predict possible scenarios in reality. On the other hand, as the existing differential equation models are built based on observations, experience, and understanding of the diseases, they often become outdated and therefore need to be updated and modified to keep up with the constant changes in epidemics. Therefore, in addition to building new models, improving existing models is also very important.

Second, once mathematical models have been formulated, aspects of qualitative study and approximate solutions are raised. Addressing these issues is useful for finding appropriate strategies for disease prevention and control, as well as for predicting disease spread scenarios. In addition, infectious diseases often need to be monitored over very long periods of time. This leads to the rapid solution of differential equation models over long time periods. Therefore, efficient numerical methods are urgently needed. However, the construction of efficient high-order numerical methods in general, and numerical methods that preserve essential qualitative properties of differential equation models in particular, is still an important problem that has not been fully solved.

Lastly, the practical application of mathematical models of infectious diseases is essential, but has not been widely used. In particular, theoretical studies should be combined with observed real-world epidemic data to calibrate the mathematical models and find optimal parameters, thereby building scenarios that better reflect reality and proposing appropriate anti-epidemic strategies.

For the above reasons, differential equation models for infectious diseases need to be studied and developed. To achieve this, it is also necessary to develop and extend research methods to keep pace with the complexity of the proposed models.

Another future direction is to use one NSFD scheme not exclusively, but as one element in a *hybrid scheme approach*, e.g. using operator splitting [11, 15], Chebyshev collocation [4], Hermite Polynomials [365] wavelets [364, 436] or a predictor corrector NSFD approach [157].

A special challenge are mimetic / fitted operator schemes for singular perturbed problems, due to the necessary resolution of boundary layers having different scales,

e.g. convection-diffusion equations [40, 41, 229], Burgers-Huxley equation [131], differential difference equations [350] or boundary value ODE problems [283, 346].

Finally, most recent research dirctions for NSFD schemes are the GPU acceleration of the (serial) NSFD code [226] and geometric numerical integration, symmetrization of NSFD schemes [85].

6. Concluding remarks and discussions

In this work, we have reviewed many but not all recent developments and real-life applications of deterministic ODEs and PDEs of major infectious diseases, mainly focusing on mathematical modeling, qualitative analysis, numerical methods and real-life applications. We have also presented and discussed some open problems and future directions that research in differential equation models for infectious diseases can take.

In the presentation, we focus only on deterministic differential equation models associated with the integer-order derivatives. Delayed models, stochastic models [14, 51, 354, 467], and fractional-order models, especially for PDEs [40, 41, 82, 418], will be considered in future work.

All the results presented demonstrate the important role of differential equation models in disease modeling. Moreover, they remain an effective and indispensable approach to study the characteristics of infectious diseases and thereby suggest effective measures for disease prevention and public health protection.

References

- M. Abdelheq, O. Belhamiti, L. Bouzid, D. Y. Trejos, J. C. Valverde, A predictive spatiotemporal model for bovine Babesiosis epidemic transmission, Journal of Theoretical Biology 480 (2019) 192-204.
- [2] E. M. Adamu, K. C. Patidar, A. Ramanantoanina, An unconditionally stable nonstandard finite difference method to solve a mathematical model describing Visceral Leishmaniasis, Mathematics and Computers in Simulation 187 (2021) 171-190.
- [3] O. Adekanye, T. Washington, Nonstandard finite difference scheme for a Tacoma Narrows Bridge model, Applied Mathematical Modelling 62 (2018) 223-236.
- [4] P. Agarwal, A. A. El-Sayed, Non-standard finite difference and Chebyshev collocation methods for solving fractional diffusion equation, Physica A: Statistical Mechanics and Its Applications 500 (2018) 40-49.
- [5] P. Agarwal, J. J. Nieto, M. Ruzhansky, D. F. M. Torres, Analysis of Infectious Disease Problems (Covid-19) and Their Global Impact, Springer, Singapore, 2021.
- [6] F. B. Agusto, A. B. Gumel, Theoretical Assessment of Avian Influenza Vaccine, Discrete and Continuous Dynamical Systems - B 13 (2010) 1-25.
- [7] F. B. Agusto, S. Lenhart, A. B. Gumel, A. Odoi, Mathematical analysis of a model for the transmission dynamics of bovine tuberculosis, Mathematical Methods in the Applied Sciences, 34 (2011) 1873-1887.
- [8] F. B. Agusto, A.B. Gumel, Qualitative dynamics of lowly- and highly-pathogenic avian influenza strains, Mathematical Biosciences 243 (2013) 147-162.
- [9] F. B. Agusto, S. Bewick, W. F. Fagan, Mathematical model of Zika virus with vertical transmission, Infectious Disease Modelling 2 (2017) 244-267.
- [10] W. Ahmad, M. Rafiq, A. I. K. Butt, N. Ahmad, T. Ismaeel, S. Malik, H. G. Rabbani, Z. Asif, Analytical and numerical explorations of optimal control techniques for the bi-modal dynamics of Covid-19, Nonlinear Dynamics (2024) 1-30.
- [11] N. Ahmed, T. S. Shaikh, M. Rafiq, M. A. Rehman, M. Ali, M. O. Ahmad, Positivity preserving operator splitting nonstandard finite difference methods for SEIR reaction diffusion model, Open Mathematics 17(1) (2019) 313-330.

- [12] N. Ahmed, M. Rafiq, D. Baleanu, A. S. Alshomrani M. A. Rehman, Positive explicit and implicit computational techniques for reaction-diffusion epidemic model of dengue disease dynamics, Advances in Difference Equations 2020 (2020), 202.
- [13] N. Ahmed, M. Rafiq, W. Adel, H. Rezazadeh, I. Khan, K. S. Nisar, Structure preserving numerical analysis of HIV and CD4+ T-cells reaction diffusion model in two space dimensions, Chaos Solitons & Fractals 139 (2020) 110307.
- [14] N. Ahmed, J. E. Macías-Díaz, A. Raza, D. Baleanu, M. Rafiq, Z. Iqbal, M. O. Ahmad, Design, analysis and comparison of a nonstandard computational method for the solution of a general stochastic fractional epidemic model, Axioms 11(1) (2021) 10.
- [15] N. Ahmed, T. S. Shaikh, M. Rafiq, S. M. Eldin, A. H. Ganie, M. Ali, A. Raza, I. Khan, M. I. Khan, Structure preserving splitting techniques for Ebola reaction-diffusion epidemic systems, Fractals 31(02) (2023) 2340041.
- [16] F. Al-Showaikh, E. Twizell, One-dimensional measles dynamics, Applied Mathematics and Computation 152 (2004), 169-194.
- [17] F. K. Alalhareth, Higher-Order Nonstandard Finite Difference Methods for Autonomous Differential Equations with Applications in Mathematical Ecology, PhD dissertation, The University of Texas at Arlington, 2022.
- [18] F. K. Alalhareth, H. V. Kojouharov, Analysis and an NSFD method of a model of bacterial competition in the presence of a plasmid, Biomath Communications Supplement (2023).
- [19] F. K. Alalhareth, M. Gupta, S. Roy, H. V. Kojouharov, Second-order modified positive and elementary stable nonstandard numerical methods for *n*-dimensional autonomous differential equations, Mathematical Methods in the Applied Sciences (2023) https://doi.org/10. 1002/mma.9560.
- [20] M. E. Alexander, C. Bowman, S. M. Moghadas, R. Summers, A. B. Gumel, B. M. Sahai, A Vaccination Model for Transmission Dynamics of Influenza, SIAM Journal on Applied Dynamical Systems 3 (2004) 503-524.
- [21] M. E. Alexander, S. M. Moghadas, Bifurcation Analysis of an SIRS Epidemic Model with Generalized Incidence, SIAM Journal on Applied Mathematics 65 (2005) 1794-1816.
- [22] W. F. Alfwzan, K. Abuasbe, A. Raza, M. Rafiq, M. Awadalla, M. A. Almulla, A nonstandard computational method for stochastic anthrax epidemic model, AIP Advances 13(7) (2023).
- [23] L. J. S. Allen, P. van den Driessche, Stochastic epidemic models with a backward bifurcation, Mathematical Biosciences and Engineering 3 (2006) 445-458.
- [24] E. Allen, Modeling with Itô Stochastic Differential Equations, Springer-Verlag, Dordrecht, The Netherlands, 2007.
- [25] L. J. S. Allen, An Introduction to Mathematical Biology, Prentice Hall, 2007.
- [26] L. J. S. Allen, B. M. Bolker, Y. Lou, A. L. Nevai, Asymptotic profiles of the steady states for an SIS epidemic reaction-diffusion model, Discrete and Continuous Dynamical Systems - B 21 (2008) 1-20.
- [27] L. J. S. Allen, An Introduction to Stochastic Epidemic Models. In: Brauer, F., van den Driessche, P., Wu, J. (eds) Mathematical Epidemiology. Lecture Notes in Mathematics, vol. 1945, 2008, Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-78911-6_3.
- [28] H. Alrabaiah, R. U. Din, K. J. Ansari, B. Ozdemir, Stability and numerical analysis via non-standard finite difference scheme of a nonlinear classical and fractional order model, Results in Physics 49 (2023) 106536.
- [29] J. Alvarez-Ramirez, F. J. Valdes-Parada, Non-standard finite-differences schemes for generalized reaction-diffusion equations, Journal of Computational and Applied Mathematics 228(1) (2009) 334-343.
- [30] R. M. Anderson, R. M. May, Infectious Diseases in Humans: Dynamics and Control, Oxford University Press, Oxford, 1991.
- [31] H. Andersson, T. Britton, Stochastic Epidemic Models and Their Statistical Analysis, Springer, New York, 2000.
- [32] C. N. Angstmann, A. M. Erickson, B. I. Henry, A. V. McGann, J. M. Murray, A. James, Fractional Order Compartment Models, SIAM Journal on Applied Mathematics 77 (2017) 430-446.
- [33] R. Anguelov, J. M.-S. Lubuma, Contributions to the mathematics of the nonstandard finite difference method and applications, Numerical Methods for Partial Differential Equations 17 (2001) 518-543.

- [34] R. Anguelov, J. M.-S. Lubuma, Nonstandard finite difference method by nonlocal approximation, Mathematics and Computers in Simulation 61 (2003) 465-475.
- [35] R. Anguelov, J. M.-S. Lubuma, S. K. Mahudu, Qualitatively stable finite difference schemes for advection-reaction equations, Journal of Computational and Applied Mathematics 158 (2003) 19-30.
- [36] R. Anguelov, P. Kama, J. M.-S. Lubuma, On non-standard finite difference models of reaction-diffusion equations, Journal of Computational and Applied Mathematics 175 (2005) 11-29.
- [37] R. Anguelov, Y. Dumont, J. M.-S. Lubum, E. Mureithi, Stability Analysis and Dynamics Preserving Nonstandard Finite Difference Schemes for a Malaria Model, Mathematical Population Studies 20 (2013) 101-122.
- [38] R. Anguelov, Y. Dumont, J. M.-S. Lubuma, M. Shillor, Dynamically consistent nonstandard finite difference schemes for epidemiological models, Journal of Computational and Applied Mathematics 255 (2014) 161-182.
- [39] R. Anguelov, T. Berge, M. Chapwanya, J. K. Djoko, P. Kama, J. M.-S. Lubuma, Y. Terefe, Nonstandard finite difference method revisited and application to the Ebola virus disease transmission dynamics, Journal of Difference Equations and Applications 26 (2020) 818-854.
- [40] W. T. Aniley, G. F. Duressa, Uniformly convergent numerical method for time-fractional convection-diffusion equation with variable coefficients, Partial Differential Equations in Applied Mathematics 8 (2023) 100592.
- [41] W. T. Aniley, G. F. Duressa, Nonstandard finite difference method for time-fractional singularly perturbed convection-diffusion problems with a delay in time. Results in Applied Mathematics 21 (2024) 100432.
- [42] S. Anita, V. Capasso, Reaction-Diffusion Systems in Epidemiology, An. Stiint. Univ. Al. I. Cuza Iasi. Mat. (N.S.) Tomul LXVI, 2020, f. 2.
- [43] A. R. Appadu, G. N. de Waal, Numerical solution of a malignant invasion model using some finite difference methods, Demonstratio Mathematica 56(1) (2023) 20220244.
- [44] D. F. Aranda, D. Y. Trejos, J. C. Valverde, R. J. Villanueva, A mathematical model for Babesiosis disease in bovine and tick populations, Mathematical Methods in the Applied Sciences 35 (2012) 249-256.
- [45] A. J. Arenas, G. González-Parra, L. Jódar, Periodic solutions of nonautonomous differential systems modeling obesity population, Chaos, Solitons & Fractals 42 (2009) 1234-1244.
- [46] A. J. Arenas, G. González-Parra, B. M. Chen-Charpentier, A nonstandard numerical scheme of predictor-corrector type for epidemic models, Computers & Mathematics with Applications 59 (2010) 3740-3749.
- [47] A. J. Arenas, G. González-Parra, B. M. Chen-Charpentier, Construction of nonstandard finite difference schemes for the SI and SIR epidemic models of fractional order, Mathematics and Computers in Simulation 121 (2016) 48-63.
- [48] J. Arino, C. C. McCluskey, P. van den Driessche, Global Results for an Epidemic Model with Vaccination that Exhibits Backward Bifurcation, SIAM Journal on Applied Mathematics 64 (2003) 260-276.
- [49] U. M. Ascher, L. R. Petzold, Computer Methods for Ordinary Differential Equations and Differential-Algebraic Equations, Society for Industrial and Applied Mathematics, Philadelphia, 1998.
- [50] E. Avila-Vales, G. E. García-Almeida, A. G. C. Pérez, Qualitative analysis of a diffusive SIR epidemic model with saturated incidence rate in a heterogeneous environment, Journal of Mathematical Analysis and Applications 503 (2021) 125295.
- [51] M. Z. Baber, A. R. Seadway, M. S. Iqbal, N. Ahmed, M. W. Yasin, M. O. Ahmed, Comparative analysis of numerical and newly constructed soliton solutions of stochastic Fisher-type equations in a sufficiently long habitat, International Journal of Modern Physics B 37(16) (2023) 2350155.
- [52] S. R. Bandekar, M. Ghosh, A co-infection model on TB-COVID-19 with optimal control and sensitivity analysis, Mathematics and Computers in Simulation 200 (2022) 1-31.
- [53] M. Banerjee, S. Ghosh, P. Manfredi, A. d'Onofrio, Spatio-temporal chaos and clustering induced by nonlocal information and vaccine hesitancy in the SIR epidemic model, Chaos, Solitons & Fractals 170 (2023) 113339.
- [54] M. Barman, N. Mishra, Hopf bifurcation analysis for a delayed nonlinear-SEIR epidemic model on networks, Chaos, Solitons & Fractals 178 (2024) 114351.

- [55] N. A. Battista, L. B. Pearcy, W. C. Strickland, Modeling the Prescription Opioid Epidemic, Bulletin of Mathematical Biology 81(2019) 2258-2289.
- [56] C. Beck, M. Hutzenthaler, A. Jentzen, B. Kuckuck, An overview on deep learning-based approximation methods for partial differential equations, Discrete and Continuous Dynamical Systems-B 28 (2023) 3697-3746.
- [57] T. Berge, J. M.-S. Lubuma, G. M. Moremedi, N. Morrisdan, R. Kondera-Shava, A simple mathematical model for Ebola in Africa, Journal of Biological Dynamics 11 (2007) 42-74.
- [58] S. Berkhahn, M. Ehrhardt, A Physics-Informed Neural Network to Model COVID-19 Infection and Hospitalization Scenarios, Advances in Continuous and Discrete Models: Theory and Applications 2022, 61 (2022).
- [59] L. Billings, I. B. Schwartz, Exciting chaos with noise: unexpected dynamics in epidemic outbreaks, Journal of Mathematical Biology 44 (2002) 31-48.
- [60] S. K. Biswas, U. Ghosh, S. Sarkar, Mathematical model of Zika virus dynamics with vector control and sensitivity analysis, Infectious Disease Modelling 5 (2020) 23-41.
- [61] S. Blanes, A. Iserles, S. Macnamara, Positivity-preserving methods for ordinary differential equations, ESAIM: M2AN 56 (2022) 1843-1870.
- [62] K. Blayneh, Y. Cao, Hee-Dae Kwon, Optimal control of vector-borne diseases: Treatment and prevention, Discrete and Continuous Dynamical Systems - B 11 (2009) 587-611.
- [63] J. Blechschmidt, O. G. Ernst, Three ways to solve partial differential equations with neural networks – A review, GAMM-Mitteilungen 44(2)(2021) e202100006.
- [64] S. M. Blower, P. M. Small, P. C. Hopewell, Control Strategies for Tuberculosis Epidemics: New Models for Old Problems, Science 273 (1996) 497-500.
- [65] S. Boccaletti, C. Grebogi, Y.-C. Lai, H. Mancini, D. Maza, The control of chaos: theory and applications, Physics Reports 329 (2000) 103-197.
- [66] B. M. Bolker, B. T. Grenfell, Chaos and biological complexity in measles dynamics, Proceedings of the Royal Society B 251 (1993) 75-81.
- [67] S. Bonhoeffer, R. M. May, G. M. Shaw, M. A. Nowak, Virus dynamics and drug therapy, The Proceedings of the National Academy of Sciences 94 (1997) 6971-6976.
- [68] F. Brauer, Backward bifurcations in simple vaccination models, Journal of Mathematical Analysis and Applications 298 (2004) 418-431.
- [69] F. Brauer, The Kermack-McKendrick epidemic model revisited, Mathematical Biosciences 198 (2005) 119-131.
- [70] F. Brauer, Compartmental Models in Epidemiology, In: Brauer, F., van den Driessche, P., Wu, J. (eds) Mathematical Epidemiology. Lecture Notes in Mathematics, vol. 1945, 2008, Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-78911-6_2.
- [71] F. Brauer, P. Driessche, J. Wu, Mathematical Epidemiology, Springer Berlin, Heidelberg, 2008.
- [72] F. Brauer, C. Castillo-Chavez, Mathematical Models in Population Biology and Epidemiology, Springer, 2012.
- [73] F. Brauer, C. Castillo-Chavez, Mathematical Models for Communicable Diseases, Society for Industrial and Applied Mathematics, Philadelphia, 2013.
- [74] F. Brauer, Mathematical epidemiology: Past, present, and future, Infectious Disease Modelling 2 (2017) 113-127.
- [75] F. Brauer, P. Driessche, Z. Feng, Mathematical Models in Epidemiology, Springer, New York, 2019.
- [76] T. Britton, Stochastic epidemic models: A survey, Mathematical Biosciences 225 (2010) 24-35.
- [77] B. Buonomo, C. Vargas-De-León, Stability and bifurcation analysis of a vector-bias model of malaria transmission, Mathematical Biosciences 142 (2013) 59-67.
- [78] B. Buonomo, D. Lacitignola, C. Vargas-De-León, Qualitative analysis and optimal control of an epidemic model with vaccination and treatment, Mathematics and Computers in Simulation 100 (2014) 88-102.
- [79] B. Buonomo, N. Chitnis, A. d'Onofrio, Seasonality in epidemic models: a literature review, Ricerche di Matematica 67 (2018) 7-25.
- [80] C. Butler, P. Stechlinski, Modeling Opioid Abuse: A Case Study of the Opioid Crisis in New England, Bulletin of Mathematical Biology 85(6) (2023) 45.

- [81] A. R. Butt, A. A. Saqib, A. S. Alshomrani, A. Bakar, M. Inc, Dynamical analysis of a nonlinear fractional cervical cancer epidemic model with the nonstandard finite difference method. Ain Shams Engineering Journal, 15(3) (2024) 102479.
- [82] L. Cai, M. Guo, Y. Li, W. Ying, H. Gao, X. Luo, Nonstandard finite difference method for nonlinear Riesz space fractional reaction-diffusion equation, International Journal of Numerical Analysis & Modeling 16(6) (2019) 925-938.
- [83] Y. Cai, S. Yan, H. Wang, X. Lian, W. Wang, Spatiotemporal Dynamics in a Reaction-Diffusion Epidemic Model with a Time-Delay in Transmission, International Journal of Bifurcation and Chaos, 25 (2015) 1550099.
- [84] Y. Cai, Y. Kang, W. Wang, A stochastic SIRS epidemic model with nonlinear incidence rate, Applied Mathematics and Computation 305 (2017) 221-240.
- [85] J. Calatayud, M. Jornet, Mathematical modeling of adulthood obesity epidemic in Spain using deterministic, frequentist and Bayesian approaches, Chaos, Solitons & Fractals 140 (2020) 110179.
- [86] J. Calatayud, M. Jornet, On the symmetrization and composition of nonstandard finite difference schemes as an alternative to Richardson's extrapolation, Journal of Difference Equations and Applications 28(5) (2022), 716-724.
- [87] W. K. Caldwell, B. Freedman, L. Settles, M. M. Thomas, E.T. Camacho, S. Wirkus, The Vicodin abuse problem: A mathematical approach, Journal of Theoretical Biology 483 (2019) 110003.
- [88] M. Calvo, M. P. Laburta, J. I. Montijano, L. Rández, Projection methods preserving Lyapunov functions, BIT Numerical Mathematics 50 (2010) 223-241.
- [89] N. Cangiotti, M. Capolli, M. Sensi, S. Sottile, A survey on Lyapunov functions for epidemic compartmental models, Bollettino dell'Unione Matematica Italiana (2023) https://doi. org/10.1007/s40574-023-00368-6.
- [90] V. Capasso, G. Serio, A generalization of the Kermack-McKendrick deterministic epidemic model, Mathematical Biosciences 42 (1978) 43-61.
- [91] V. Capasso, Reaction-Diffusion Models for the Spread of a Class of Infectious Diseases. In: Neunzert, H. (eds) Proceedings of the Second European Symposium on Mathematics in Industry. European Consortium for Mathematics in Industry, vol. 3, 1988, Springer, Dordrecht. https://doi.org/10.1007/978-94-009-2979-1_11.
- [92] A. M. Carvalho, S. Goncalves, An analytical solution for the Kermack–McKendrick model, Physica A 566 (2021) 125659.
- [93] R. Casagrandi, L. Bolzoni, S. A. Levin, V. Andreasen, The SIRC model and influenza A, Mathematical Biosciences 200 (2006) 152-169.
- [94] M. A. Castro, C. J. Mayorga, A. Sirvent, F. Rodríguez, Exact numerical solutions and high order nonstandard difference schemes for a second order delay differential equation, Mathematical Methods in the Applied Sciences 46(17) (2023) 17962-17979.
- [95] I. T. Cetinkaya, An Application of Nonstandard Finite Difference Method to a Model Describing Diabetes Mellitus and Its Complications, Journal of New Theory 45 (2023) 105-119.
- [96] A. Champneys, K. Tsaneva-Atanasova, Dynamical Systems Theory, Bifurcation Analysis, Encyclopedia of Systems Biology, pp. 632-637.
- [97] K. Chang, Z. Zhang, G. Liang, Dynamics analysis of a nonlocal diffusion dengue model, Scientific Reports 13(1) (2023) 15239.
- [98] L. Chang, W. Gong, Z. Jin, G.-Q. Sun, Sparse Optimal Control of Pattern Formations for an SIR Reaction-Diffusion Epidemic Model, SIAM Journal on Applied Mathematics 82(2022) 1764-1790.
- [99] L. Chang, X. Wang, G. Sun, Z. Wang, Z. Jin, A time independent least squares algorithm for parameter identification of Turing patterns in reaction-diffusion systems, Journal of Mathematical Biology 88(1) (2024) 5.
- [100] M. Chapwanya, J. M.-S. Lubuma, R. E. Mickens, From enzyme kinetics to epidemiological models with Michaelis-Menten contact rate: Design of nonstandard finite difference schemes, Computers & Mathematics with Applications 64 (2012) 201-213. equation; nonstandard finite difference metho
- [101] M. Chapwanya, J. M.-S. Lubuma, R. E. Mickens, Nonstandard finite difference schemes for Michaelis–Menten type reaction-diffusion equations, Numerical Methods for Partial Differential Equations 29(1) (2013) 337-360.

- [102] M. Chapwanya, J. M.-S. Lubuma, R. E. Mickens, Positivity-preserving nonstandard finite difference schemes for cross-diffusion equations in biosciences, Computers & Mathematics with Applications 68 (2014) 1071-1082.
- [103] L. Chen, J. Sun, Optimal vaccination and treatment of an epidemic network model, Physics Letters A 378 (2014) 3028-3036.
- [104] W.-Y. Chen, S. Bokka, Stochastic modeling of nonlinear epidemiology, Journal of Theoretical Biology 234(4) (2005) 455-470.
- [105] Y. Chen, F. Liu, Q. Yu, T. Li, Review of fractional epidemic models, Applied Mathematical Modelling 97 (2021) 281-307.
- [106] B. M. Chen-Charpentier, D. T. Dimitrov, H. V. Kojouharov, Combined nonstandard numerical methods for ODEs with polynomial right-hand sides, Mathematics and Computers in Simulation 73 (2006) 105-113.
- [107] B. M. Chen-Charpentier, H. V. Kojouharov, An unconditionally positivity preserving scheme for advection-diffusion reaction equations, Mathematical and Computer Modelling 57(9-10) (2013) 2177-2185.
- [108] C. Cheng, Z. Zheng, Dynamics and spreading speed of a reaction-diffusion system with advection modeling West Nile virus, Journal of Mathematical Analysis and Applications 493 (2021) 124507.
- [109] N. Chitnis, J. M. Cushing, J. M. Hyman, Bifurcation Analysis of a Mathematical Model for Malaria Transmission, SIAM Journal on Applied Mathematics 67 (2006) 24-45.
- [110] G. Chowell, P. W. Fenimore, M. A. Castillo-Garsow, C. Castillo-Chavez, SARS outbreaks in Ontario, Hong Kong and Singapore: the role of diagnosis and isolation as a control mechanism, Journal of Theoretical Biology 224 (2003) 1-8.
- [111] G. Chowell, Fitting dynamic models to epidemic outbreaks with quantified uncertainty: A primer for parameter uncertainty, identifiability, and forecasts, Infectious Disease Modelling 2 (2017) 379-398.
- [112] D. P. Clemence-Mkhope, Dynamically Consistent NSFD Discretization of Some Productive-Destructive Population Models Satisfying Conservations Laws, Open Access Library Journal 8(4) (2021) 1-12.
- [113] S. Cole, S. Wirkus, Modeling the Dynamics of Heroin and Illicit Opioid Use Disorder, Treatment, and Recovery, Bulletin of Mathematical Biology 84 (2022) 48.
- [114] D. Conte, G. Pagano, B. Paternoster, Nonstandard finite differences numerical methods for a vegetation reaction-diffusion model, Journal of Computational and Applied Mathematics 419 (2023) 114790.
- [115] K. L. Cooke, P. van den Driessche, Analysis of an SEIRS epidemic model with two delays, Journal of Mathematical Biology 35 (1996) 240-260.
- [116] A. Coronel, F. Huancas, I. Hess, A. Tello, The diffusion identification in a SIS reactiondiffusion system, Mathematical Biosciences and Engineering 21 (2024) 562-581.
- [117] C. Cosner, J. C. Beier, R. S. Cantrell, D. Impoinvil, L. Kapitanski, M. D. Potts, A. Troyo, S. Ruan, The effects of human movement on the persistence of vector-borne diseases, Journal of Theoretical Biology 258 (2009) 550-560.
- [118] G. M. R. Costa, M. Lobosco, M. Ehrhardt, R. F. Reis, Mathematical Analysis and a Nonstandard Scheme for a Model of the Immune Response against COVID-19, in: A. Gumel (ed.), Mathematical and Computational Modeling of Phenomena Arising in Population Biology and Nonlinear Oscillations: In honour of the 80th birthday of Ronald E. Mickens, AMS Contemporary Mathematics, 2023.
- [119] J. Cresson, F. Pierret, Non standard finite difference scheme preserving dynamical properties, Journal of Computational and Applied Mathematics 303 (2016) 15-30.
- [120] J. Cresson, A. Szafrańska, Discrete and continuous fractional persistence problems the positivity property and applications, Communications in Nonlinear Science and Numerical Simulation 44 (2017) 424-448.
- [121] N. Crokidakis, L. Sigaud, Modeling the evolution of drinking behavior: A Statistical Physics perspective, Physica A 570 (2021) 125814.
- [122] R. Cui, King-Yeung Lam, Y. Lou, Dynamics and asymptotic profiles of steady states of an epidemic model in advective environments, Journal of Differential Equations 263 (2017) 2343-2373.
- [123] Q. A. Dang, M. T. Hoang, Dynamically consistent discrete metapopulation model, Journal of Difference Equations and Applications 22 (2016) 1325-1349.

- [124] Q. A. Dang, M. T. Hoang, Lyapunov direct method for investigating stability of nonstandard finite difference schemes for metapopulation models, Journal of Difference Equations and Applications 24 (2018) 15-47.
- [125] Q. A. Dang, M. T. Hoang, Nonstandard finite difference schemes for a general predator-prey system, Journal of Computational Science 36 (2019) 101015.
- [126] Q. A. Dang, M. T. Hoang, Positive and elementary stable explicit nonstandard Runge-Kutta methods for a class of autonomous dynamical systems, International Journal of Computer Mathematics 97 (2020) 2036-2054.
- [127] Q. A. Dang, M. T. Hoang, Positivity and global stability preserving NSFD schemes for a mixing propagation model of computer viruses, Journal of Computational and Applied Mathematics 374 (2020) 112753.
- [128] Q. A. Dang, M. T. Hoang, Exact finite difference schemes for three dimensional linear systems with constant coefficient, Vietnam Journal of Mathematics 46 (2018) 471–492.
- [129] M. Z. Dauhoo, B. S. N. Korimboccus, S. B. Issack, On the dynamics of illicit drug consumption in a given population, IMA Journal of Applied Mathematics 78 (2013) 432-448.
- [130] A. Dénes, A. B. Gumel, Modeling the impact of quarantine during an outbreak of Ebola virus disease, Infectious Disease Modelling 4 (2019) 12-27.
- [131] E. B. Derzie, J. B. Munyakazi, T. G. Dinka, A NSFD method for the singularly perturbed Burgers-Huxley equation, Frontiers in Applied Mathematics and Statistics 9 (2023) 1068890.
- [132] G. N. de Waal, A. R. Appadu, C. J. Pretorius, Some standard and nonstandard finite difference schemes for a reaction-diffusion-chemotaxis model, Open Physics 21(1) (2023) 20220231.
- [133] O. Diekmann, J. A. P. Heesterbeek, J. A. J. Metz, On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations, Journal of Mathematical Biology 28 (1990) 365-382.
- [134] K. Dietz, Overall patterns in the transmission cycle of infectious disease agents, In: R.M. Anderson, R.M. May (eds.) Population Biology of Infectious Diseases. Life Sciences Research Report, Vol. 25. Springer, Berlin Heidelberg New York, pp. 87-102 (1982).
- [135] D. T. Dimitrov, H. V. Kojouharov, Complete mathematical analysis of predator-prey models with linear prey growth and Beddington-DeAngelis functional response, Applied Mathematics and Computation 162 (2005) 523-538.
- [136] D. T. Dimitrov, H. V. Kojouharov, Nonstandard finite-difference schemes for general twodimensional autonomous dynamical systems, Applied Mathematics Letters 18 (2005) 769-774.
- [137] D. T. Dimitrov, H. V. Kojouharov, Positive and elementary stable nonstandard numerical methods with applications to predator-prey models Journal of Computational and Applied Mathematics 189(1-2) (2006) 98-108.
- [138] D. T. Dimitrov, H. V. Kojouharov, Stability-Preserving Finite-Difference Methods for General Multi-Dimensional Autonomous Dynamical Systems, International Journal of Numerical Analysis and Modeling 4(2) (2007) 282-292.
- [139] D. T. Dimitrov, H. V. Kojouharov, Nonstandard finite-difference methods for predator-prey models with general functional response, Mathematics and Computers in Simulation 78(1) (2008) 1-11.
- [140] D. T. Dimitrov, H. V. Kojouharov, Dynamically consistent numerical methods for general productive-destructive systems, Journal of Difference Equations and Applications 17 (2011) 1721-1736.
- [141] R. U. Din, K. A. Khan, A. Aloqaily, N. Mlaiki, H. Alrabaiah, Using Non-Standard Finite Difference Scheme to Study Classical and Fractional Order SEIVR Model, Fractal and Fractional 7(7) (2023) 552.
- [142] D. Ding, X. Ding, Dynamic consistent non-standard numerical scheme for a dengue disease transmission model, Journal of Difference Equations and Applications 20(3) (2014) 492-505.
- [143] A. d'Onofrio, P. Manfredi, E. Salinelli, Vaccinating behaviour, information, and the dynamics of SIR vaccine preventable diseases, Theoretical Population Biology 71 (2007) 301-317.
- [144] A. d'Onofrio, On pulse vaccination strategy in the SIR epidemic model with vertical transmission, Applied Mathematics Letters 18 (2005) 729-732.
- [145] X. Duana S. Yuan, Z. Qiu, J. Ma, Global stability of an SVEIR epidemic model with ages of vaccination and latency, Computers & Mathematics with Applications 68 (2014) 288-308.

- [146] J. Dushoff, W. Huang, C. Castillo-Chavez, Backwards bifurcations and catastrophe in simple models of fatal diseases, Journal of Mathematical Biology 36 (1998) 227-248.
- [147] D. J. D. Earn, P. Rohani, B. T. Grenfell, Persistence, chaos and synchrony in ecology and epidemiology, Proceedings of the Royal Society of London. Series B 26:57-10.
- [148] M. Ehrhardt, R. E. Mickens, A nonstandard finite difference scheme for convection-diffusion equations having constant coefficients, Applied Mathematics and Computation 219 (2013) 6591-6604.
- [149] M. Ehrhardt, J. Gašper, S. Kilianová, SIR-based Mathematical Modeling of Infectious Diseases with Vaccination and Waning Immunity, Journal of Computational Science 37 (2019) 101027.
- [150] K. Ejima, K. Aihara, H. Nishiura, Modeling the obesity epidemic: social contagion and its implications for control, Theoretical Biology and Medical Modelling 10 (2013), 1-13.
- [151] A. M. Elaiw, N. H. AlShamrani, Global stability of humoral immunity virus dynamics models with nonlinear infection rate and removal, Nonlinear Analysis: Real World Applications 26 (2015) 161-190.
- [152] A. M. Elaiw, M. Alshaikh, Stability preserving NSFD scheme for a general virus dynamics model with antibody and cell-mediated responses, Chaos Solitons & Fractals 138 (2020) 109862.
- [153] E. H. Elbasha, C. N. Podder, A. B. Gumel, Analyzing the dynamics of an SIRS vaccination model with waning natural and vaccine-induced immunity, Nonlinear Analysis: Real World Applications 12 (2011) 2692-2705.
- [154] E. H. Elbasha, Model for hepatitis C virus transmissions, Mathematical Biosciences and Engineering 10 (2013) 1045-1065.
- [155] S. Elsheikh, R. Ouifki, K. C. Patidar, A non-standard finite difference method to solve a model of HIV-malaria co-infection, Journal of Difference Equations and Applications 20(3) (2014) 354-378.
- [156] I. Faragó, R. Mosleh, Some qualitative properties of the discrete models for malaria propagation, Applied Mathematics and Computation 438 (2023) 127628.
- [157] A. Farooqi, R. Ahmad, H. Alotaibi, T. A. Nofal, R. Farooqi, I. Khan, A comparative epidemiological stability analysis of predictor corrector type non-standard finite difference scheme for the transmissibility of measles, Results in Physics 21 (2021) 103756.
- [158] H. Fatoorehchi, M. Ehrhardt, Numerical and semi-numerical solutions of a modified Thévenin model for calculating terminal voltage of battery cells, Journal of Energy Storage 145 (2022) 103746.
- [159] Z. Feng, H. R. Thieme, Endemic Models with Arbitrarily Distributed Periods of Infection I: Fundamental Properties of the Model, SIAM Journal on Applied Mathematics 61 (2000) 803 - 833.
- [160] Z. Feng, H. R. Thieme, Endemic Models with Arbitrarily Distributed Periods of Infection II: Fast Disease Dynamics and Permanent Recovery, SIAM Journal on Applied Mathematics 61 (2000) 983-1012.
- [161] F. Forouzanni, A. B. Gumel, Mathematical analysis of an age-structured model for malaria transmission dynamics, Mathematical Biosciences 247 (2014) 80-94.
- [162] A. F. Fossi, J. Lubuma, C. Tadmon, B. Tsanou, Mathematical modeling and nonstandard finite difference scheme analysis for the environmental and spillover transmissions of Avian Influenza A model, Dynamical Systems 36 (2021) 212-255.
- [163] B. M. Fundzama, Design, analysis and simulation of a robust numerical method to solve Zika virus models, Master Thesis, University of the Western Cape, 2019. https://etd.uwc. ac.za/handle/11394/7122
- [164] C. Gai, D. Iron, T. Kolokolnikov, Localized outbreaks in an S-I-R model with diffusion, Journal of Mathematical Biology 80 (2020) 1389-1411.
- [165] C. Gan, X. Yang, W. Liu, Q. Zhu, A propagation model of computer virus with nonlinear vaccination probability, Communications in Nonlinear Science and Numerical Simulation 19 (2014) 92-100.
- [166] S. M. Garba, A. B. Gumel, J.-S. Lubuma, Dynamically-consistent non-standard finite difference method for an epidemic model, Mathematical and Computer Modelling 53(1–2) (2011) 131–150.

- [167] C. C. Garsow, G. J. Salivia, A. R. Herrera, Mathematical Models for the Dynamics of Tobacco use, recovery and relapse, Technical Report Series BU-1505-M, Cornell University, UK (2000).
- [168] Y. Geng, J. Xu, Global stability of a delayed and diffusive virus model with nonlinear infection function, Journal of Biological Dynamics 15 (2021) 287-307.
- [169] A. Gerisch, R. Weiner, The positivity of low-order explicit Runge-Kutta schemes applied in splitting methods, Computers & Mathematics with Applications 45 (2003) 53-67.
- [170] P. Glendinning, L. P. Perry, Melnikov analysis of chaos in a simple epidemiological model, Journal of Mathematical Biology 35 (1997) 359-373.
- [171] G. González-Parra, A. J. Arenas, B. M. Chen-Charpentier, Combination of nonstandard schemes and Richardson's extrapolation to improve the numerical solution of population models, Mathematical and Computer Modelling 52 (2010) 1030-1036.
- [172] E. González, M. J. Villena, On the spatial dynamics of vaccination: A spatial SIRS-V model, Computers & Mathematics with Applications 80 (2020) 733-743.
- [173] A. Gragnani, S. Rinaldi, G. Feichtinger, Dynamics of drug consumption: a theoretical model, Socio-Economic Planning Sciences 31 (1997) 127-137.
- [174] A. Gray, D. Greenhalgh, L. Hu, X. Mao, J. Pan, A Stochastic Differential Equation SIS Epidemic Model, SIAM Journal on Applied Mathematics 71 (2011) 876-902.
- [175] B. T. Grenfell, B. M. Bolker, A. Kleczkowski, Seasonality and extinction in chaotic metapopulations, Proc. Proceedings of the Royal Society of London. Series B 259:97-103.
- [176] V. Grimm, G. R. W. Quispel, Geometric Integration Methods that Preserve Lyapunov Functions, BIT Numerical Mathematics 45 (2005) 709-723.
- [177] A. B. Gumel, S. M. Moghadas, A qualitative study of a vaccination model with non-linear incidence, Applied Mathematics and Computation 143 (2003) 409-419.
- [178] A. B. Gumel, R. E. Mickens, B. D. Corbett, A non-standard finite-difference scheme for a model of HIV transmission and control, Journal of Computational Methods in Sciences and Engineering 3(1) (2003) 91-98.
- [179] A. B. Gumel, S. Ruan, T. Day, J. Watmough, F. Brauer, P. van den Driessche, D. Gabrielson, Ch. Bowman, M. E. Alexander, S. Ardal, J. Wu, B. M. Sahai, Modelling strategies for controlling SARS outbreaks, Proceedings of the Royal Society of London B 271 (2004) 2223-2232.
- [180] A. B. Gumel, C. C. McCluskey, J. Watmough, An SVEIR Model for Assessing Potential Impact of an Imperfect Anti-SARS Vaccine, Mathematical Biosciences 3 (2006) 485-512.
- [181] A. B. Gumel, B. Song, Existence of multiple-stable equilibria for a multi-drug-resistant model of mycobacterium tuberculosis, Mathematical Biosciences and Engineering 5 (2008) 437-455.
- [182] A. B. Gumel, Global dynamics of a two-strain avian influenza model, International Journal of Computer Mathematics 86 (2009) 85-108.
- [183] A. B. Gumel, Causes of backward bifurcations in some epidemiological models, Journal of Mathematical Analysis and Applications 395 (2012) 355-365.
- [184] K. F. Gurski, A simple construction of nonstandard finite-difference schemes for small nonlinear systems applied to SIR models, Computers & Mathematics with Applications 66 (2013) 2165-2177.
- [185] K. P. Hadeler, P. Van den Driessche, Backward bifurcation in epidemic control, Mathematical Biosciences 146 (1997) 15-35.
- [186] E. Hairer, G. Wanner, S. P. Norsett, Solving Ordinary Differential Equations I: Nonstiff Problems, Springer Berlin, Heidelberg, 1993.
- [187] E. Hairer, G. Wanner, Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems, Springer Berlin, Heidelberg, 1996.
- [188] E. Hairer, Ch. Lubich, G. Wanner, Geometric Numerical Integration, Springer-Verlag, 2002.
- [189] T. Harko, F. S. N. Lobo, M. K. Mak, Exact analytical solutions of the Susceptible-Infected-Recovered (SIR) epidemic model and of the SIR model with equal death and birth rates, Applied Mathematics and Computation 236 (2014) 184-194.
- [190] J. A. P. Heesterbeek, J. A. J. Metz, The saturating contact rate in marriage and epidemic models, Journal of Mathematical Biology 31 (1993) 529-539.
- [191] F. Heldmann, S. Berkhahn, M. Ehrhardt, K. Klamroth, PINN Training using Biobjective Optimization: The Trade-off between Data Loss and Residual Loss, Journal of Computational Physics 488 (2023), 112211.

- [192] S. Henshaw, C. Connell McCluskey, Global stability of a vaccination model with immigration, Electronic Journal of Differential Equations 2015 (2015) 1-10.
- [193] E. Hernandez-Martinez, H. Puebla, F. Valdes-Parada, J. Alvarez-Ramirez, Nonstandard finite difference schemes based on Green's function formulations for reaction-diffusionconvection systems, Chemical Engineering Science 94 (2013) 245-255.
- [194] J. E. Herrera-Serrano, J. E. Macías-Díaz, I. E. Medina-Ramírez, J. A. Guerrero, An efficient nonstandard computer method to solve a compartmental epidemiological model for COVID-19 with vaccination and population migration, Computer Methods and Programs in Biomedicine 221 (2022) 106920.
- [195] H. W. Hethcote, M. A. Lewis, P. van den Driessche, An epidemiological model with a delay and a nonlinear incidence rate, Journal of Mathematical Biology 27 (1989) 49-64
- [196] H. W. Hethcote, The mathematics of Infectious diseases, SIAM Review 42 (2000) 599-653.
- [197] S. Hews, S. Eikenberry, J. D. Nagy, Y. Kuang, Rich dynamics of a hepatitis B viral infection model with logistic hepatocyte growth, Journal of Mathematical Biology 60 (2010) 573-590.
- [198] M. W. Hirsch, S. Smale, R. L. Devaney, Differential Equations, Dynamical Systems, and an Introduction to Chaos, Third Edition, Elsevier, 2013.
- [199] M. T. Hoang, Dynamically consistent nonstandard finite difference schemes for a virus-patch dynamic model, Journal of Applied Mathematics and Computing 68 (2022) 3397-3423.
- [200] M. T. Hoang, A novel second-order nonstandard finite difference method for solving onedimensional autonomous dynamical systems, Communications in Nonlinear Science and Numerical Simulation 114 (2022) 106654.
- [201] M. T. Hoang, Reliable approximations for a hepatitis B virus model by nonstandard numerical schemes, Mathematics and Computers in Simulation 193(2022) 32-56.
- [202] M. T. Hoang, Dynamical analysis of a generalized hepatitis B epidemic model and its dynamically consistent discrete model, Mathematics and Computers in Simulation 205 (2023) 291-314.
- [203] M. T. Hoang, M. Ehrhardt, A dynamically consistent nonstandard finite difference scheme for a generalized SEIR epidemic model, Journal of Difference Equations and Applications (Dec. 2023) https://doi.org/10.1080/10236198.2023.2291151.
- [204] M. T. Hoang, A class of second-order and dynamically consistent nonstandard finite difference schemes for nonlinear Volterra's population growth model, Computational and Applied Mathematics 42 (2023) Article number: 85.
- [205] M. T. Hoang, A novel second-order nonstandard finite difference method preserving dynamical properties of a general single-species model, International Journal of Computer Mathematics 100 (2023) 2047-2062.
- [206] M. T. Hoang, J. C. Valverde, A generalized model for the population dynamics of a two stage species with recruitment and capture using a nonstandard finite difference scheme, Computational and Applied Mathematics 43 (2024) article no. 54.
- [207] M. T. Hoang, M. Ehrhardt, A general class of second-order L-stable explicit numerical methods for stiff problems, Applied Mathematics Letters 149 (2024) 108897.
- [208] M. T. Hoang, M. Ehrhardt, A second-order nonstandard finite difference method for a general Rosenzweig-MacArthur predator-prey model, Journal of Computational and Applied Mathematics 444(2024) 115752.
- [209] Z. Horváth, Positivity of Runge-Kutta and diagonally split Runge-Kutta methods, Applied Numerical Mathematics 28 (1998) 309-326.
- [210] S.-B. Hsu, Lih-Ing W. Roeger, The final size of a SARS epidemic model without quarantine, Journal of Mathematical Analysis and Applications 333 (2007) 557-566.
- [211] Y. Hu, Q. Pan, W. Hou, M. He, Rumor spreading model considering the proportion of wisemen in the crowd, Physica A 505 (2018) 1084-1094.
- [212] Z. Hu, W. Ma, S. Ruan, Analysis of SIR epidemic models with nonlinear incidence rate and treatment, Mathematical Biosciences 238 (2012) 12-20.
- [213] W. Huang, M. Han, K. Liu, Dynamics of an SIS reaction-diffusion epidemic model for disease transmission, Mathematical Biosciences and Engineering 7 (2010) 51-66.
- [214] H.-F. Huo, H. Xue, H. Xiang, Dynamics of an alcoholism model on complex networks with community structure and voluntary drinking, Physica A 505 (2018) 880-890.
- [215] M. Iannelli, M. Martcheva, X.-Z. Li, Strain replacement in an epidemic model with superinfection and perfect vaccination, Mathematical Biosciences 195 (2005) 23-46.

- [216] M. A. Ibrahim, A. Dénes, Threshold and stability results in a periodic model for malaria transmission with partial immunity in humans, Applied Mathematics and Computation 392 (2021) 125711.
- [217] A. Iggidr, G. Sallet, M. O. Souza, On the dynamics of a class of multi-group models for vector-borne diseases, Journal of Mathematical Analysis and Applications 441 (2016) 723-743.
- [218] Z. Iqbal, J. E. Macías-Díaz, N. Ahmed, M. A.-U. Rehman, A. Raza, M. Rafiq, A SEIR model with memory effects for the propagation of Ebola-like infections and its dynamically consistent approximation, Computer Methods and Programs in Biomedicine 209 (2021) 106322.
- [219] Z. Iqbal, M. A.-U. Rehman, M. Imran, N. Ahmed, U. Fatima, A. Akgül, M. Rafiq, A. Raza, A. A. Djuraev, F. Jarad, A finite difference scheme to solve a fractional order epidemic model of computer virus, AIMS Mathematics 8 (2023) 2337-2359.
- [220] J. Jang, H.-D. Kwon, J. Lee, Optimal control problem of an SIR reaction-diffusion model with inequality constraints, Mathematics and Computers in Simulation 171 (2020) 136-151.
- [221] M. Jawaz, N. Ahmed, D. Baleanu, M. Rafiq, M. A. Rehman, Positivity preserving technique for the solution of HIV/AIDS reaction diffusion model with time delay, Frontiers in Physics 7 (2020) 229.
- [222] Y. Jin, W. Wang, S. Xiao, An SIRS model with a nonlinear incidence rate, Chaos, Solitons & Fractals 34 (2007) 1482-1497.
- [223] L. Jódar, F.-J. Santonja, G. Gonzalez-Párra, Modelling dynamics of infant obesity in the region of Valencia, Spain, Computers & Mathematics with Applications 56 (2008) 679-689.
- [224] L. Joáar, R. J. Villanueva, A. J. Arenas, Gilberto C. González, Nonstandard numerical methods for a mathematical model for influenza disease, Mathematics and Computers in Simulation 79 (2008) 622-633.
- [225] A. Jones, N. Strigu, Is spread of COVID-19 a chaotic epidemic, Chaos, Solitons & Fractals 142 (2021) 110376.
- [226] Y. Kanai, T. Hoshino, T. Ohtani, N. V. Kantartzis, GPU Acceleration of the Nonstandard FDTD Method, In: 2023 International Applied Computational Electromagnetics Society Symposium (ACES), pp. IEEE, 2023, pp. 1-2.
- [227] S. Kayenat, A. K. Verma, NSFD schemes for a class of nonlinear generalised advection-diffusion-reaction equation, Pramana 96(1) (2022) 14.
- [228] L. Kalachev, E. L. Landguth, J. Graham, Revisiting classical SIR modelling in light of the COVID-19 pandemic, Infectious Disease Modelling 8 (2023) 72-83.
- [229] O. O. Kehinde, J. B. Munyakazi A. R. Appadu, A NSFD Discretization of Two-Dimensional Singularly Perturbed Semilinear Convection-Diffusion Problems, Frontiers in Applied Mathematics and Statistics 8 (2022) 861276.
- [230] W. O. Kermack, A. G. McKendrick, A contribution to the mathematical theory of epidemics, Proceedings of the Royal Society of London - Series A 115 (1927) 700-721.
- [231] W. O. Kermack, A. G. McKendrick, Contributions to the mathematical theory of epidemics. II. -The problem of endemicity, Proceedings of the Royal Society of London - Series A 138 (1932) 55-83.
- [232] W. O. Kermack, A. G. McKendrick, Contributions to the mathematical theory of epidemics. III. -Further studies of the problem of endemicity, Proceedings of the Royal Society of London - Series A 141 (1933) 94-122.
- [233] P. G. Kevrekidis, J. Cuevas-Maraver, Y. Drossinos, Z. Rapti, G. A. Kevrekidis, Reactiondiffusion spatial modeling of COVID-19: Greece and Andalusia as case examples, Physical Review E 104 (2021) 024412.
- [234] H. K. Khalil, Nonlinear systems, Prentice Hall, 2002.
- [235] M. M. Khalsaraei, A positive and elementary stable nonstandard explicit scheme for a mathematical model of the influenza disease, Mathematics and Computers in Simulation 182 (2021) 397-410.
- [236] M. A. Khan, S. Islam, J. C Valverde, S. A. Khan, Control strategies of hepatitis B with three control variables, Journal of Biological Systems26(2018) 1-21.
- [237] H. Khan, R. N. Mohapatra, K. Vajravelu, S.J. Liao, The explicit series solution of SIR and SIS epidemic models, Applied Mathematics and Computation 215 (2009) 653-669.

- [238] I. U. Khan, A. Hussain, S. Li, A. Shokri, Modeling the Transmission Dynamics of Coronavirus Using Nonstandard Finite Difference Scheme, Fractal and Fractional 7(6) (2023) 451.
- [239] M. A. Khan, S. W. Shah, S. Ullah, J. F. Gomez-Aguilar, A dynamical model of asymptomatic carrier zika virus with optimal control strategies, Nonlinear Analysis: Real World Applications 50 (2019) 144-170.
- [240] M. I. Khan, K. Al-Khaled, A. Raza, S. U. Khan, J. Omar, A. M. Galal, Mathematical and numerical model for the malaria transmission: Euler method scheme for a malarial model, International Journal of Modern Physics B 37(16) (2023) 2350158.
- [241] O. Khyar, K. Allali, Global dynamics of a multi-strain SEIR epidemic model with general incidence rates: application to COVID-19 pandemic, Nonlinear Dynamics 102 (2020) 489-509.
- [242] K. I. Kim, Z. Lin, L. Zhang, Avian-human influenza epidemic model with diffusion, Nonlinear Analysis: Real World Applications 11 (2010) 313-322.
- [243] K. Kitagawa , S. Nakaok, Y. Asai, K. Watashi, S. Iwami, A PDE multiscale model of hepatitis C virus infection can be transformed to a system of ODEs, Journal of Theoretical Biology 448 (2018) 80-85.
- [244] F. Köhler-Rieper, C. H. F. Röhl, E. De Micheli, A novel deterministic forecast model for the Covid-19 epidemic based on a single ordinary integro-differential equation, The European Physical Journal Plus 135(7) (2020) 599.
- [245] H. V. Kojouharov, S. Roy, M. Gupta, F. Alalhareth, J. M. Slezak, A second-order modified nonstandard theta method for one-dimensional autonomous differential equations, Applied Mathematics Letters 112 (2021) 106775.
- [246] S. Kopecz, A. Meister, On order conditions for modified Patankar-Runge-Kutta schemes, Applied Numerical Mathematics 123 (2018) 159-179.
- [247] S. Kopecz, A. Meister, On the existence of three-stage third-order modified Patankar-Runge-Kutta schemes, Numerical Algorithms 81 (2019) 1473-1484.
- [248] A. Korobeinikov, Lyapunov functions and global properties for SEIR and SEIS epidemic models, Mathematical Medicine and Biology 21 (2004) 75-83.
- [249] A. Korobeinikov, P. K. Maini, A Lyapunov function and global properties for SIR and SEIR epidemiological models with nonlinear incidence, Mathematical Biosciences & Engineering 1 (2004) 57-60.
- [250] A. Korobeinikov, Lyapunov Functions and Global Stability for SIR and SIRS Epidemiological Models with Non-Linear Transmission, Bulletin of Mathematical Biology 68 (2006) 615-626.
- [251] T. Kossaczka, M. Ehrhardt, M. Günther, Deep FDM: Enhanced finite difference methods by deep learning, Franklin Open 4 (2023) 100039.
- [252] E. Kovács, J. Majár, M. Saleh, Unconditionally Positive, Explicit, Fourth Order Method for the Diffusion-and Nagumo-Type Diffusion-Reaction Equations, Journal of Scientific Computing, 98(2) (2024) 39.
- [253] C. M. Kribs-Zaleta, J. X. Velasco-Hernández, A simple vaccination model with multiple endemic states, Mathematical Biosciences 164 (2000) 183-201.
- [254] N. A. Kudryashov, M. A. Chmykhov, M. Vigdorowitsch, Analytical features of the SIR model and their applications to COVID-19, Applied Mathematical Modelling 90 (2021) 466-473.
- [255] T. Kuniya, J. Wang, Lyapunov functions and global stability for a spatially diffusive SIR epidemic model, Applicable Analysis 96 (2017) 1935-1960.
- [256] T. Kuniya, J. Wang, Global dynamics of an SIR epidemic model with nonlocal diffusion, Nonlinear Analysis: Real World Applications 43 (2018) 262-282.
- [257] Y. A. Kuznetsov, C. Piccardi, Bifurcation analysis of periodic SEIR and SIR epidemic models, Journal of Mathematical Biology 32 (1994) 109-121.
- [258] Y. A. Kuznetsov, Elements of Applied Bifurcation Theory, Springer New York, 2004.
- [259] W. Lacarbonara, J. Ma, C. Nataraj, Preface to the special issue "Complex dynamics of COVID-19: modeling, prediction and control (part II)", Nonlinear Dynamics 109 (2022) 1-3.
- [260] A. Lahrouz, L. Omari, D. Kiouach, A. Belmaâti, Deterministic and stochastic stability of a mathematical model of smoking, Statistics and Probability Letters 81 (2011) 1276-1284.

- [261] A. Lahrouz, L. Omari, D. Kiouach, A. Belmaâti, Complete global stability for an SIRS epidemic model with generalized non-linear incidence and vaccination, Applied Mathematics and Computation 218 (2012) 6519-6525.
- [262] J. La Salle, S. Lefschetz, Stability by Liapunov's Direct Method. Academic Press, New York, 1961.
- [263] A. A. Lashari, G. Zaman, Global dynamics of vector-borne diseases with horizontal transmission in host population, Computers & Mathematics with Applications 61 (2011) 745-754.
- [264] J. Legrand, R. F. Grais, P. Y. Boelle, A. J. Valleron, A. Flahault, Understanding the dynamics of Ebola epidemics, Epidemiology & Infection 135 (2007) 610-621.
- [265] R. J. LeVeque, Finite Difference Methods for Ordinary and Partial Differential Equations, Society for Industrial and Applied Mathematics, Philadelphia, 2007.
- [266] B. Li, Q. Bie, Long-time dynamics of an SIRS reaction-diffusion epidemic model, Journal of Mathematical Analysis and Applications 475 (2019) 1910-1926.
- [267] C.-L. Li, C.-H. Li, C.-Y. Cheng, Analysis of an epidemiological model with age of infection, vaccination, quarantine and asymptomatic transmission, Journal of the Franklin Institute 360 (2023) 657-692.
- [268] J. Li, H. Jiang, X. Mei, C. Hu, G. Zhang, Dynamical analysis of rumor spreading model in multi-lingual environment and heterogeneous complex networks, Information Sciences 536 (2020) 391-408.
- [269] M. Y. Li, J. S. Muldowney, On Bendixson's criterion, Journal of Differential Equations 106 (1993) 27-39.
- [270] M. Y. Li, J. S. Muldowney, Global stability for the SEIR model in epidemiology, Mathematical Biosciences 125 (1995) 155-164.
- [271] M. Y. Li, J. S. Muldowney, A geometric approach to global-stability problems, SIAM Journal on Mathematical Analysis 27 (1996) 1070-1083.
- [272] M. Y. Li, J. S. Muldowney, P. van den Driessche, Global stability of SEIRS models in epidemiology, Canadian Applied Mathematics Quarterly 7(4) (1999).
- [273] S. Li, I. Bukhsh, I. U. Khan, M. I. Asjad, S. M. Eldin, M. A. El-Rahman, D. Baleanu The impact of standard and nonstandard finite difference schemes on HIV nonlinear dynamical model, Chaos, Solitons & Fractals 173 (2023) 113755.
- [274] X. Li, R. P. Agarwal, J. F. Gomez-Aguilar, Q. Badshah, G. ur Rahman, Threshold dynamics: Formulation, stability & sensitivity analysis of co-abuse model of heroin and smoking, Chaos, Solitons & Fractals 161 (2022) 112373.
- [275] Y. Li, X. Liu, Y. Yuan, J. Li, L. Wang, Global analysis of tuberculosis dynamical model and optimal control strategies based on case data in the United States, Applied Mathematics and Computation 422 (2022) 126983.
- [276] H.-l. Lin, F.-B. Wang, Global dynamics of a nonlocal reaction-diffusion system modeling the West Nile virus transmission, Nonlinear Analysis: Real World Applications 46 (2019) 352-373.
- [277] C. Liu, R. Cui, Qualitative analysis on an SIRS reaction-diffusion epidemic model with saturation infection mechanism, Nonlinear Analysis: Real World Applications 62 (2021) 103364.
- [278] P. Liu, S. N. Elaydi, Discrete Competitive and Cooperative Models of Lotka-Volterra Type, Journal of Computational Analysis and Applications 3 (2001) 53-73.
- [279] X. Liu, B. Dai, Flip bifurcations of an SIR epidemic model with birth pulse and pulse vaccination, Applied Mathematical Modelling 43 (2017) 579-591.
- [280] X.-L. Liu, C.-C. Zhu, A non-standard finite difference scheme for a diffusive HIV-1 infection model with immune response and intracellular delay, Axioms 11(3) (2022) 129.
- [281] Y. Lou, R. B. Salako, Mathematical analysis of the dynamics of some reaction-diffusion models for infectious diseases, Journal of Differential Equations 370 (2023) 424-469.
- [282] X. Lu, H. Hui, F. Liu, Y. Bai, Stability and optimal control strategies for a novel epidemic model of COVID-19, Nonlinear Dynamics 106 (2021) 1491-150.
- [283] J. M.-S. Lubuma, K. C. Patidar, Uniformly convergent non-standard finite difference methods for self-adjoint singular perturbation problems, Journal of Computational and Applied Mathematics 191(2) (2006) 228–238.
- [284] A. M. Lyapunov, The General Problem of the Stability of Motion. Taylor & Francis, London, 1992.

- [285] M. H. Maamar, M. Ehrhardt, L. Tabharit, A nonstandard finite difference scheme for a timefractional model of Zika virus transmission, Mathematical Biosciences and Engineering 21(1) (2024) 924-962.
- [286] P. Magal, G. Webb, The parameter identification problem for SIR epidemic models: identifying unreported cases, Journal of Mathematical Biology 77 (2018) 1629-1648.
- [287] P. Magal, G. F. Webb, Y. Wu, On the Basic Reproduction Number of Reaction-Diffusion Epidemic Models, SIAM Journal on Applied Mathematics 79 (2019) 284-304.
- [288] Y. Mammeri, A reaction-diffusion system to better comprehend the unlockdown: Application of SEIR-type model with diffusion to the spatial spread of COVID-19 in France, Computational and Mathematical Biophysics 8(1) (2020) 102-113.
- [289] S. Mandal, R. R. Sarkar, S. Sinha, Mathematical models of malaria a review, Malaria Journal 2011, 10:202.
- [290] S. Mangiarotti, Low dimensional chaotic models for the plague epidemic in Bombay (1896-1911), Chaos, Solitons & Fractals 81 (2015) 184-196.
- [291] S. Mangiarotti, M. Peyre, M. Huc, A chaotic model for the epidemic of Ebola virus disease in West Africa (2013-2016), Chaos 26 (2016) 113112.
- [292] S. Mangiarotti, M. Peyre, Y. Zhang, M. Huc, F. Roger, Y. Kerr, Chaos theory applied to the outbreak of COVID-19: an ancillary approach to decision making in pandemic context, Epidemiology and Infection 148 (2020), e95, 1-9.
- [293] K. Manna, S. P. Chakrabarty, Global stability and a non-standard finite difference scheme for a diffusion driven HBV model with capsids, Journal of Difference Equations and Applications 21 (2015) 918-933.
- [294] K. Manna, A non-standard finite difference scheme for a diffusive HBV infection model with capsids and time delay, Journal of Difference Equations and Applications 23 (2017) 1901-1911.
- [295] B. P. Marchant, J. Norbury, A. J. Perumpanani, Traveling shock waves arising in a model of malignant invasion, SIAM Journal on Applied Mathematics 60(2) (2000) 463-476.
- [296] T. T. Marinov, R. S. Marinova, Inverse problem for adaptive SIR model: Application to COVID-19 in Latin America, Infectious Disease Modelling 7 (2022) 134-148.
- [297] M. Martcheva, An Introduction to Mathematical Epidemiology, Springer New York, NY, 2015.
- [298] N. K. Martin, P. Vickerman, M. Hickman, Mathematical modelling of hepatitis C treatment for injecting drug users, Journal of Theoretical Biology 274 (2011) 58-66.
- [299] J. Martin-Vaquero, A. Martin del Rey, A. H. Encinas, J. D. Hernandez Guillen, A. Queiruga-Dios, G. Rodriguez Sanchez, Higher-order nonstandard finite difference schemes for a MSEIR model for a malware propagation, Journal of Computational and Applied Mathematics 317 (2017) 146-156.
- [300] J. Martin-Vaquero, A. Queiruga-Dios, A. Martin del Rey, A. H. Encinas, J. D. Hernandez Guillen, G. Rodriguez Sanchez, Variable step length algorithms with high-order extrapolated non-standard finite difference schemes for a SEIR model, Journal of Computational and Applied Mathematics 330 (2018) 848-854.
- [301] J. P. S. Maurício de Carvalho, A. A. Rodrigues, SIR Model with Vaccination: Bifurcation Analysis, Qualitative Theory of Dynamical Systems 22 (2023) article number 105.
- [302] R. M. May, Chaos and the dynamics of biological populations, Proceedings of the Royal Society A 413(1844) (1987) 27-44.
- [303] C. C. McCluskey, Y. Yang, Global stability of a diffusive virus dynamics model with general incidence function and time delay, Nonlinear Analysis: Real World Applications 25 (2015) 64-78.
- [304] A. McNabb, Comparison theorems for differential equations, Journal of Mathematical Analysis and Applications 119 (1986) 417-428.
- [305] M. Mehdizadeh Khalsaraei, R. Shokri Jahandizi, Positivity-preserving nonstandard finite difference schemes for simulation of advection-diffusion reaction equations, Computational Methods for Differential Equations 2(4) (2014) 256-267.
- [306] J. Mena-Lorca, H.W. Hethcote, Dynamic models of infectious diseases as regulators of population size, Journal of Mathematical Biology 30 (1992) 693-716.
- [307] R. E. Mickens, I. Ramadhani, Finite-difference schemes having the correct linear stability properties for all finite step-sizes III, Computers & Mathematics with Applications 27 (1994) 77-84.

- [308] R. E. Mickens, Nonstandard Finite Difference Models of Differential Equations, World Scientific, Singapore, 1994.
- [309] R. E. Mickens, Nonstandard finite difference schemes for reaction-diffusion equation, Numerical Methods for Partial Differential Equation 15 (1999) 201-214.
- [310] R. E. Mickens, Nonstandard finite difference schemes for reaction-diffusion equations having linear advection, Numerical Methods for Partial Differential Equation 16 (2000) 361-364.
- [311] R. E. Mickens, Applications of Nonstandard Finite Difference Schemes, World Scientific, Singapore, 2000.
- [312] R. E. Mickens, A nonstandard finite-difference scheme for the Lotka-Volterra system, Applied Numerical Mathematics 45 (2003) 309-314
- [313] R. E. Mickens, A nonstandard finite difference scheme for a Fisher PDE having nonlinear diffusion, Computers & Mathematics with Applications 45(1-3) (2003) 429-436.
- [314] R. E. Mickens, A nonstandard finite difference scheme for a PDE modeling combustion with nonlinear advection and diffusion, Mathematics and Computers in Simulation 69 (2005) 439-446.
- [315] R. E. Mickens, Dynamic consistency: a fundamental principle for constructing nonstandard finite difference schemes for differential equations, Journal of Difference Equations and Applications 11 (2005) 645-653.
- [316] R. E. Mickens, Advances in the Applications of Nonstandard Finite Difference Schemes, World Scientific, Singapore, 2005.
- [317] R. E. Mickens, Numerical integration of population models satisfying conservation laws: NSFD methods, Journal of Biological Dynamics 4 (2007) 427-436.
- [318] R. E. Mickens, Calculation of denominator functions for nonstandard finite difference schemes for differential equations satisfying a positivity condition, Numerical Methods for Partial Differential Equation 23 (2007) 672-691.
- [319] R. E. Mickens, Nonstandard Finite Difference Schemes for Differential Equations, Journal of Difference Equations and Applications 8 (2012) 823-847.
- [320] R. E. Mickens, T. M. Washington, NSFD discretizations of interacting population models satisfying conservation laws, Computers & Mathematics with Applications 66 (2013) 2307-2316.
- [321] R. E. Mickens, Nonstandard Finite Difference Schemes: Methodology and Applications, World Scientific, 2020.
- [322] R. E. Mickens, T. M. Washington, A note on a positivity preserving nonstandard finite difference scheme for a modified parabolic reaction-advection-diffusion PDE, Journal of Difference Equations and Applications 26 (2020) 1423-1427.
- [323] R. E. Mickens, I. H. Herron, Approximate rational solutions to the Thomas-Fermi equation based on dynamic consistency, Applied Mathematics Letters 116 (2021) 106994.
- [324] L. Min, Y. Su, Y. Kuang, Mathematical analysis of a basic virus infection model with application to HBV infection, Rocky Mountain Journal of Mathematics 38 (2008), 1573-1585.
- [325] T. Y. Miyaoka, S. Lenhart, J. F. C. A. Meyer, Optimal control of vaccination in a vectorborne reaction-diffusion model applied to Zika virus, Journal of Mathematical Biology 79 (2019) 1077-1104.
- [326] N. Mohan, N. Kumari, Positive steady states of a SI epidemic model with cross diffusion, Applied Mathematics and Computation 410 (2021) 126423.
- [327] J. Mondal, S. Khajanchi, Mathematical modeling and optimal intervention strategies of the COVID-19 outbreak, Nonlinear Dynamics 109 (2022) 177-202.
- [328] G. Mulone, B. Straughan, A note on heroin epidemics, Mathematical Biosciences 218 (2009) 138-141.
- [329] A. Mummert, O. M. Otunuga, Parameter identification for a stochastic SEIRS epidemic model: case study influenza, Journal of Mathematical Biology 79 (2019) 705-729.
- [330] L. H. A. Monteiro, An epidemiological model for SARS-CoV-2, Ecological Complexity 43 (2020) 100836.
- [331] G. A. Munoz-Fernandez, J. M. Seoane, J. B. Seoane-Sepulveda, A SIR-type model describing the successive waves of COVID-19, Chaos, Solitons & Fractals 144 (2021) 110682.
- [332] J. G. Murray, Mathematical Biology, II: Spatial Models and Biomedical Applications, third ed., Springer, New York, 2003.

- [333] E. M. Musyoki, W. N. Mutuku, N. M. Imbusi, E. O. Omondi, Mathematical modelling of tuberculosis and diabetes co-infection using the non-standard finite difference scheme, Pan-American Journal of Mathematics 2 (2023) 16.
- [334] M. C. Nava, J. M. Guevara-Jordan, A New Analysis of an Implicit Mimetic Scheme for the Heat Equation, Journal of Applied Mathematics and Physics 11(3) (2023) 841-857.
- [335] F. Ndairou, M. Khalighi, L. Lahti, Ebola epidemic model with dynamic population and memory, Chaos, Solitons & Fractals 170 (2023) 113361.
- [336] M. Z. Ndii, A. K. Supriatna, An application of nonstandard finite-difference scheme for solving autonomous and non-autonomous mathematical model for Wolbachia-carrying mosquito population dynamics, Journal of Mathematical and Computational Science 11(1) (2021) 1039-1052.
- [337] T. W. Ng, G. Turinici, A. Danchin, A double epidemic model for the SARS propagation, BMC Infectious Diseases 3 (2003) 19.
- [338] G. A. Ngwa, On the Population Dynamics of the Malaria Vector, Bulletin of Mathematical Biology 68 (2006) 2161-2189.
- [339] M. A. Nowak, S. Bonhoeffer, A. M. Hill, R. Boehme R, H. C. Thomas, H. McDade, Viral dynamics in hepatitis B virus infection, The Proceedings of the National Academy of Sciences 93 (1996) 4398-4402.
- [340] M. A. Nowak, R. M. May, Virus Dynamics: Mathematical Principles of immunology and Virology, Oxford University Press, New York, 2000.
- [341] H. A. Obaid, R. Ouifki, K. C. Patidar, A nonstandard finite difference method for solving a mathematical model of HIV-TB co-infection, Journal of Difference Equations and Applications 23(6) (2017) 1105-1132.
- [342] L. F. Olsen, G. L. Truty, W. M. Schaffer, Oscillations and chaos in epidemics: A nonlinear dynamic study of six childhood diseases in Copenhagen, Denmark, Theoretical Population Biology 33 (1988) 344-370.
- [343] L. F. Olsen, W. M. Schaffer, Chaos Versus Noisy Periodicity: Alternative Hypotheses for Childhood Epidemics, Science 249 (1990) 499-505.
- [344] S. M. O'Regan, T. C. Kelly, A. Korobeinikov, M. J. A. O'Callaghan, A. V. Pokrovskii, Lyapunov functions for SIR and SIRS epidemic models, Applied Mathematics Letters 23 (2010) 446-448.
- [345] R. Padmanabhan, H. S. Abed, N. Meskin, T. Khattab, M. Shraim, M. A. Al-Hitmi, A review of mathematical model-based scenario analysis and interventions for COVID-19, Computer Methods and Programs in Biomedicine 209 (2021) 106301.
- [346] P. K. Pandey, A method for an approximate numerical solution of two point boundary value problems: nonstandard finite difference method on semi open interval, International Journal of Computing Science and Mathematics 17(3) (2023) 220-228.
- [347] S. A. Pasha, Y. Nawaz, M. S. Arif, On the nonstandard finite difference method for reactiondiffusion models, Chaos, Solitons & Fractals 166 (2023) 112929.
- [348] M. Pathak, P. Joshi, K. S. Nisar, Numerical investigation of fluid flow and heat transfer in micropolar fluids over a stretching domain, Journal of Thermal Analysis and Calorimetry 147 (2022) 10637-10646.
- [349] K. C. Patidar, On the use of nonstandard finite difference methods, Journal of Difference Equations and Applications 11 (2005) 735-758.
- [350] K. C. Patidar, K. K. Sharma, ε-Uniformly convergent non-standard finite difference methods for singularly perturbed differential difference equations with small delay, Applied Mathematics and Computation 175(1) (2006) 864–890.
- [351] K. C. Patidar, Nonstandard finite difference methods: Recent trends and further developments, Journal of Difference Equations and Applications 22 (2016) 817-849.
- [352] A. Perasso, B. Laroche, Y. Chitour, S. Touzeau, Identifiability analysis of an epidemiological model in a structured population, Journal of Mathematical Analysis and Applications 374 (2011) 154-165.
- [353] V. Piccirillo, Nonlinear control of infection spread based on a deterministic SEIR model, Chaos, Solitons & Fractals 149 (2021) 111051.
- [354] F. Pierret, A non-standard-Euler-Maruyama scheme, Journal of Difference Equations and Applications 22(1) (2016) 75–98.
- [355] J. R. C. Piqueira, O. V. Araujo, A modified epidemiological model for computer viruses, Applied Mathematics and Computation 213 (2009) 355-360.

- [356] J. R. C. Piqueira, M. Zilbovicius, C. M. Batistela, Daley-Kendal models in fake-news scenario, Physica A 548 (2020) 123406.
- [357] L. Pontryagin, V. Boltyanskii, R. Gramkrelidze, E. Mischenko, The Mathematical Theory of Optimal Processes, Wiley Interscience, 1962.
- [358] W. Qin, L. Wang, X. Ding, A non-standard finite difference method for a hepatitis B virus infection model with spatial diffusion, Journal of Difference Equations and Applications 20 (2014) 1641-1651.
- [359] W. Qin, D. Ding, X. Ding, A non-standard finite difference scheme for an advection-diffusionreaction equation, Mathematical Methods in the Applied Sciences 38(15) (2015) 3308-3321.
- [360] M. Rafiq, M. O. Ahmad, Non-Standard Finite Difference Modeling for Transmission Dynamics of Dengue Fever, University of Engineering and Technology Taxila. Technical Journal 21(1) (2016) 116.
- [361] M. Rafiq, J. E. Macías-Díaz, A. Raza, N. Ahmed, Design of a nonlinear model for the propagation of COVID-19 and its efficient nonstandard computational implementation, Applied Mathematical Modelling 89 (2021) 1835-1846.
- [362] M. Rahman, K. Bekele-Maxwell, L. L. Cates, H. T. Banks, N. K. Vaidya, Modeling Zika Virus transmission Dynamics: parameter estimates, Disease Characteristics, and prevention, Scientific Reports 9 (2019) 10575.
- [363] M. Raissi, P. Perdikaris, G. E. Karniadakis, Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations, Journal of Computational Physics 378 (2019) 686-707.
- [364] M. K. Rawani, A. K. Verma, C. Cattani, A novel hybrid approach for computing numerical solution of the time-fractional nonlinear one and two-dimensional partial integrodifferential equation, Communications in Nonlinear Science and Numerical Simulation 118 (2023) 106986.
- [365] M. K. Rawani, A. K. Verma, L. Verma, Numerical treatment of Burgers' equation based on weakly L-stable generalized time integration formula with the NSFD scheme, Applied Mathematics and Computation 467 (2024) 128485.
- [366] A. Raza, M. S. Arif, M. Rafiq, A reliable numerical analysis for stochastic dengue epidemic model with incubation period of virus, Advances in Difference Equations 2019(1) (2019) 1-19.
- [367] A. Raza, U. Fatima, M. Rafiq, N. Ahmed, I. Khan, K. S. Nisar, Z. Iqbal, Mathematical analysis and design of the nonstandard computational method for an epidemic model of computer virus with delay effect: Application of mathematical biology in computer science, Results in Physics 21 (2021) 103750.
- [368] J. Ren, X. Yang, Q. Zhu, L.-X. Yang, C. Zhang, A novel computer virus model and its dynamics, Nonlinear Analysis: Real World Applications 13 (2012) 376-384.
- [369] X. Ren, K. Wang, X. Liu, Dynamics on a degenerated reaction-diffusion Zika transmission model, Applied Mathematics Letters 150 (2024) 108935.
- [370] M. Riaz, K. Shah, A. Ullah, M. A. Alqudah, T. Abdeljawad, The Volterr-Lyapunov matrix theory and nonstandard finite difference scheme to study a dynamical system, Results in Physics 52 (2023) 106890.
- [371] F. A. Rihan, Delay differential equations and applications to biology, Springer, Singapore, 2021.
- [372] P. Rodrigues, M. Gabriela, M. Gomes, C. Rebelo, Drug resistance in tuberculosis-a reinfection model, Theoretical Population Biology 71 (2007) 196-212.
- [373] L.-I. W. Roeger, Dynamically consistent discrete Lotka-Volterra competition models derived from nonstandard finite-difference schemes, Discrete and Continuous Dynamical Systems -Series B 9(2) (2008) 415-429.
- [374] L.-I. W. Roeger, G. Lahodny Jr., Dynamically consistent discrete Lotka-Volterra competition systems, Journal of Difference Equations and Applications 19 (2013) 191-200.
- [375] G. Rohith, K. B. Devik, Dynamics and control of COVID-19 pandemic with nonlinear incidence rates, Nonlinear Dynamics 101 (2020) 2013-2026.
- [376] S. Ruan, W. Wang, Dynamical behavior of an epidemic model with a nonlinear incidence rate, Journal Differential Equations 188 (2003) 135-163.
- [377] M. A. Safi, A. B. Gumel, Mathematical analysis of a disease transmission model with quarantine, isolation and an imperfect vaccine, Computers & Mathematics with Applications 61 (2011) 3044-3070.

- [378] P. Saha, N. Bairagi, G. N'Guerekata, Positivity and dynamics preserving discretization schemes for nonlinear evolution equations, Malaya Journal of Matematik 12(01) (2024) 1-20.
- [379] S. M. Salman, A nonstandard finite difference scheme and optimal control for an HIV model with Beddington–DeAngelis incidence and cure rate, The European Physical Journal Plus 135 (2020) 1-23.
- [380] Md. Samsuzzoha, M. Singh, D. Lucy, Parameter estimation of influenza epidemic model, Applied Mathematics and Computation 220 (2013) 616-629.
- [381] F. Sanchez, M. Engman, L. Harrington, and C. Castillo-Chavez, Models for dengue transmission and control. In: Mathematical Studies on Human Disease Dynamics: Emerging Paradigms and Challenges. American Mathematical Society Contemporary Mathematics Series, A. B. Gumel, C. Castillo-Chavez, R. E. Mickens, D. P. Clemence (Eds.), 2007.
- [382] F.-J. Santonja, E. Sánchez, M. Rubio J.-M. Morera, Alcohol consumption in Spain and its economic cost: A mathematical modelling approach, Mathematical and Computer Modelling 52 (2010), 999-1003.
- [383] F.-J. Santonja, I.-C. Lombana, M. Rubio, E. Sánchez, J. Villanueva, A network model for the short-term prediction of the evolution of cocaine consumption in Spain, Mathematical and Computer Modelling 52 (2010) 1023-1029.
- [384] F.-J. Santonja, R.-J. Villanueva, L. Jódar, G. Gonzalez-Parra, Mathematical modelling of social obesity epidemic in the region of Valencia, Spain, Mathematical and Computer Modelling of Dynamical Systems 16 (2010) 23-34.
- [385] F.-J. Santonja, A. C. Tarazona, R. J. Villanueva, Predicting Cocaine Consumption in Spain. A mathematical Modelling Approach, Drugs: Education, Prevention & Policy 18 (2011), 108-115.
- [386] T. Sarkar, O. Biswas, P. K. Srivastava, Modelling the effects of media information and saturated treatment on malaria disease with NSFD method, International Journal of Biomathematics (2024).
- [387] W. E. Schiesser, A Mathematical Modeling Approach to Infectious Diseases: Cross Diffusion PDE Models for Epidemiology, World Scientific Publishing, 2018.
- [388] E. Scholl, H. G. Schuster, Handbook of Chaos Control, Wiley-VCH, 2008.
- [389] N. C. Severo, Generalizations of some stochastic epidemic models, Mathematical Biosciences 4 (1969) 395-402.
- [390] N. Shahid, M. A. Rehman, A. Khalid, U. Fatima, T. S. Shaikh, N. Ahmed, H. Alotaibi, M. Rafiq, I. Khan, K. S. Nisar, Mathematical analysis and numerical investigation of advection-reaction-diffusion computer virus model, Results in Physics 26 (2021) 104294.
- [391] C. Shan, H. Zhu, Bifurcations and complex dynamics of an SIR model with the impact of the number of hospital beds, Journal of Differential Equations 257 (2014) 1662-1688.
- [392] N. Sharma, R. Singh, J. Singh, O. Castillo, Modeling assumptions, optimal control strategies and mitigation through vaccination to Zika virus, Chaos, Solitons & Fractals 150 (2021) 111137.
- [393] O. Sharomi, A. B. Gumel, Curtailing smoking dynamics: A mathematical modeling approach, Applied Mathematics and Computation 195 (2008) 475-499.
- [394] O. Sharomi, C. N. Podder, A. B. Gumel, S. M. Mahmud, E. Rubinstein, Modelling the Transmission Dynamics and Control of the Novel 2009 Swine Influenza (H1N1) Pandemic, Bulletin of Mathematical Biology 73 (2011) 515-548.
- [395] O. Sharomi, T. Malik, Optimal control in epidemiology, Annals of Operations Research 251 (2017) 55-71. (2017)
- [396] L. Shi, Z. Chen, P. Wu, Spatial and temporal dynamics of COVID-19 with nonlocal dispersal in heterogeneous environment: Modeling, analysis and simulation, Chaos, Solitons & Fractals 174 (2023) 113891.
- [397] Z. Shuai, P. van den Driessche, Global Stability of Infectious Disease Models Using Lyapunov Functions, SIAM Journal on Applied Mathematics 73 (2013) 1513 - 1532.
- [398] B. Shulgin, L. Stone, Z. Agur, Pulse vaccination strategy in the SIR epidemic model, Bulletin of Mathematical Biology 60 (1998) 1123-1148.
- [399] R. P. Sigdel, C. C. McCluskey, Global stability for an SEI model of infectious disease with immigration, Applied Mathematics and Computation 243 (2014) 684-689.

- [400] M. Sinan, K. J. Ansari, A. Kanwal, K. Shah, T. Abdeljawad, Zakirullah, B. Abdalla, Analysis of the mathematical model of cutaneous Leishmaniasis disease, Alexandria Engineering Journal 72 (2023) 117-134.
- [401] R. A. Singh, R. Lal, R. R. Kotti, Time-discrete SIR model for COVID-19 in Fiji, Epidemiology and Infection 150, e75, 1-10.
- [402] G. D. Smith, Numerical Solution of Partial Differential Equations: Finite Difference Methods, Third Edition, Oxford University Press, 1985.
- [403] H. Smith, P. Waltman, The Theory of the Chemostat: Dynamics of Microbial Competition, Cambridge University Press, 1995.
- [404] H. Smith, An Introduction to Delay Differential Equations with Applications to the Life Sciences, Springer, New York, 2011.
- [405] B. Song, C. Castillo-Chavez, J. P. Aparicio, Tuberculosis models with fast and slow dynamics: the role of close and casual contacts, Mathematical Biosciences 180 (2002) 187-205.
- [406] P. Song, Y. Lou, Y. Xiao, A spatial SEIRS reaction-diffusion model in heterogeneous environment, Journal of Differential Equations 267 (2019) 5084-5114.
- [407] Y. Song, T. Zhang, Y. Peng, Turing-Hopf bifurcation in the reaction-diffusion equations and its applications, Communications in Nonlinear Science and Numerical Simulation 33 (2016) 229-258.
- [408] M. E. Songolo, B. Bidégaray-Fesquet, Extending nonstandard finite difference scheme rules to systems of nonlinear ODEs with constant coefficients, Journal of Difference Equations and Applications (2024) 1-26.
- [409] S. Soubeyrand, L. Roques, Parameter estimation for reaction-diffusion models of biological invasions, Population Ecology 56 (2014) 427-434.
- [410] P. T. Sowndarrajan, L. Shangerganesh, A. Debbouche, D. F. M. Torres, Optimal control of a heroin epidemic mathematical model, Optimization: A Journal of Mathematical Programming and Operations Research 71 (2022) 3107-3131.
- [411] T. Stocksa, L. J. Martin, S. Kuhlmann-Berenzon, T. Britto, Dynamic modeling of hepatitis C transmission among people who inject drugs, Epidemics 30 (2020) 100378.
- [412] J. C. Strikwerda, Finite Difference Schemes and Partial Differential Equations, Second Edition, Society for Industrial and Applied Mathematics, Philadelphia, 2004.
- [413] A. Stuart, A. R. Humphries, Dynamical Systems and Numerical Analysis, Cambridge University Press, 1998.
- [414] H. Sun, Y. Zhang, D. Baleanu, W. Chen, Y. Chen, A new collection of real world applications of fractional calculus in science and engineering, Communications in Nonlinear Science and Numerical Simulation 64 (2018) 213-231.
- [415] C. Tadmon, S. Foko, Modeling and mathematical analysis of an initial boundary value problem for hepatitis B virus infection, Journal of Mathematical Analysis and Applications 474 (2019) 309-350.
- [416] C. Tadmon, S. Foko, Non-standard finite difference method applied to an initial boundary value problem describing hepatitis B virus infection, Journal of Difference Equations and Applications 26 (2020) 122-139.
- [417] C. Tadmon, J. N. Kengne, Mathematical modelling and nonstandard finite scheme analysis for an Ebola model transmission with information and voluntary isolation, Journal of Difference Equations and Applications 28 (2022) 299-334.
- [418] M. Taghipour, H. Aminikhah, An Efficient Non-standard Finite Difference Scheme for Solving Distributed Order Time Fractional Reaction–Diffusion Equation, International Journal of Applied and Computational Mathematics 8(2) (2022) 56.
- [419] B. Takács, R. Horváth, I. Faragó, Space dependent models for studying the spread of some diseases, Computers & Mathematics with Applications 80 (2020) 395-404.
- [420] S. Y. Tchoumi, M. L. Diagne, H. Rwezaura, J. M. Tchuenche, Malaria and COVID-19 codynamics: A mathematical model and optimal control, Applied Mathematical Modelling 99 (2021) 294-327.
- [421] Y. A. Terefe, H. Gaff, M. Kamga, L. van der Mescht, Mathematics of a model for Zika transmission dynamics, Theory in Biosciences 137 (2018) 209-218.
- [422] J. J. Tewa, S. Bowong, S. C. O. Noutchie, Mathematical analysis of a two-patch model of tuberculosis disease with staged progression, Applied Mathematical Modelling 36 (2012) 5792-5807.

- [423] J. W. Thomas, Numerical Partial Differential Equations: Finite Difference Methods, Springer New York, 1995.
- [424] Y. Tian, X. Liu, Global dynamics of a virus dynamical model with general incidence rate and cure rat, Nonlinear Analysis: Real World Applications 16 (2014) 17-26.
- [425] S. Treibert, H. Brunner, M. Ehrhardt, Compartment models for vaccine effectiveness and non-specific effects for Tuberculosis, Mathematical Biosciences and Engineering 16(6) (2019), 7250-7298.
- [426] S. Treibert, H. Brunner, M. Ehrhardt, A nonstandard finite difference scheme for the SVICDR model to predict COVID-19 dynamics, Mathematical Biosciences and Engineering 19 (2022) 1213-1238.
- [427] D. Y. Trejos, J. C. Valverde, E. Venturino, Dynamics of infectious diseases: A review of the main biological aspects and their mathematical translation, Applied Mathematics and Nonlinear Sciences 7 (2022) 1-26.
- [428] J. Tumwiine, J. Y. T. Mugisha, L. S. Luboobi, On global stability of the intra-host dynamics of malaria and the immune system, Journal of Mathematical Analysis and Applications 341 (2008) 855-869.
- [429] P. van den Driessche, Some epidemiological models with delays, In: Differential Equations and Applications to Biology and to Industry (Claremont, CA, 1994), pages 507-520. World Sci. Publishing, River Edge, NJ (1996).
- [430] P. van den Driessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, Mathematical Biosciences 180 (2002) 29-48.
- [431] P. van den Driessche, J. Watmough, Further Notes on the Basic Reproduction Number. In: Brauer, F., van den Driessche, P., Wu, J. (eds) Mathematical Epidemiology. Lecture Notes in Mathematics, vol. 1945, 2008, Springer, Berlin, Heidelberg. https://doi.org/10.1007/ 978-3-540-78911-6_6.
- [432] P. van den Driessche, Reproduction numbers of infectious disease models, Infectious Disease Modelling 2 (2017) 288-303.
- [433] G. A. K. van Voorn, B. W. Kooi, Smoking epidemic eradication in a eco-epidemiological dynamical model, Ecological Complexity 14 (2013) 180-189.
- [434] S. Vaz, Sandra, D. F. M. Torres, A dynamically-consistent nonstandard finite difference scheme for the SICA model, arXiv preprint arXiv:2105.10826 (2021).
- [435] C. Vargas-De-León, On the global stability of SIS, SIR and SIRS epidemic models with standard incidence, Chaos, Solitons & Fractals 44 (2011) 1106-1110.
- [436] A. K. Verma, M. K. Rawani, Numerical solutions of generalized Rosenau-KDV-RLW equation by using Haar wavelet collocation approach coupled with nonstandard finite difference scheme and quasilinearization, Numerical Methods for Partial Differential Equations 39(2) (2023) 1085-1107.
- [437] A. Viguerie, A. Veneziani, G. Lorenzo, D. Baroli, N. Aretz-Nellesen, A. Patton, T. E. Yankeelov, A. Reali, T. J. R. Hughes, F. Auricchio, Diffusion-reaction compartmental models formulated in a continuum mechanics framework: application to COVID-19, mathematical analysis, and numerical study, Computational Mechanics 66 (2020) 1131-1152.
- [438] B. Wacker, J. C. Schluter, Time-continuous and time-discrete SIR models revisited: theory and applications, Advances in Difference Equations 2020 (2020) 556.
- [439] B. Wacker, J. C. Schluter, A non-standard finite-difference-method for a non-autonomous epidemiological model: analysis, parameter identification and applications, Mathematical Biosciences and Engineering 20(7) (2023) 12923-12954.
- [440] G. D. Walters, Spontaneous remission from alcohol, tobacco, and other drug abuse: seeking quantitative answers to qualitative questions, The American Journal of Drug and Alcohol Abuse 26 (2000), 443-460.
- [441] C. Wan, T. Li, W. Zhang, J. Dong, Dynamics of epidemic spreading model with drugresistant variation on scale-free networks, Physica A 493 (2018) 17-28.
- [442] J. Wang, R. Zhang, T. Kuniya, A reaction-diffusion Susceptible-Vaccinated-Infected-Recovered model in a spatially heterogeneous environment with Dirichlet boundary condition, Mathematics and Computers in Simulation 190 (2021) 848-865.
- [443] J. Wang, X. Wu, T. Kuniya, Analysis of a diffusive HBV model with logistic proliferation and non-cytopathic antiviral mechanisms, Communications in Nonlinear Science and Numerical Simulation 106 (2022) 106110.

- [444] J. Wang, Z. Teng, B. Dai, Qualitative analysis of a reaction-diffusion SIRS epidemic model with nonlinear incidence rate and partial immunity, Infectious Disease Modelling 8 (2023) 881-911.
- [445] K. Wang, W. Wang, Propagation of HBV with spatial dependence, Mathematical Biosciences 210 (2007) 78-95.
- [446] K. Wang, A. Fan, A. Torres, Global properties of an improved hepatitis B virus model, Nonlinear Analysis: Real World Applications 11 (2010) 3131-3138.
- [447] L. Wang, L.-I. W. Roeger, Nonstandard finite difference schemes for a class of generalized convection-diffusion-reaction equations, Numerical Methods for Partial Differential Equations 31 (2015) 1288-1309.
- [448] L. Wang, H. Zhao, Modeling and dynamics analysis of Zika transmission with contaminated aquatic environments, Nonlinear Dynamics 104 (2021) 845-862.
- [449] W. Wang, X.-Q. Zhao, Basic Reproduction Numbers for Reaction-Diffusion Epidemic Models, SIAM Journal on Applied Dynamical Systems 11 (2012) 1652-1673.
- [450] H. Wei, X. Li, M. Martcheva, An epidemic model of a vector-borne disease with direct transmission and time delay, Journal of Mathematical Analysis and Applications 342 (2008) 895-908.
- [451] E. White, C. Comiskey, Heroin epidemics, treatment and ODE modeling, Mathematical Biosciences 208 (2007), 312-324.
- [452] D. T. Wood, D. T. Dimitrov, H. V. Kojouharov, A nonstandard finite difference method for n-dimensional productive-destructive systems, Journal of Difference Equations and Applications 21 (2015) 240-254.
- [453] D. T. Wood, H. V. Kojouharov, A class of nonstandard numerical methods for autonomous dynamical systems, Applied Mathematics Letters 50 (2015) 78-82.
- [454] D. T. Wood, H. V. Kojouharov, D. T. Dimitrov, Universal approaches to approximate biological systems with nonstandard finite difference methods, Mathematics and Computers in Simulation 133 (2017) 337-350.
- [455] P. Wu, X. Wang, Z. Feng, Spatial and temporal dynamics of SARS-CoV-2: Modeling, analysis and simulation, Applied Mathematical Modelling 113 (2023) 220-240.
- [456] H. Xiang, Ying-Ping Liu, Hai-Feng Huo, Stability of an SAIRS alcoholism model on scalefree networks, Physica A 473 (2017) 276-292.
- [457] M.-Z. Xin, B.-G. Wang, Global dynamics of a reaction-diffusion malaria model, Nonlinear Analysis: Real World Applications 61 (2021) 103332.
- [458] J. Xu, Y. Geng, J. Hou, A non-standard finite difference scheme for a delayed and diffusive viral infection model with general nonlinear incidence rate, Computers & Mathematics with Applications 74(8) (2017) 1782–1798.
- [459] R. Xu, Z. Ma, An HBV model with diffusion and time delay, Journal of Theoretical Biology 257 (2009) 499-509.
- [460] J. Yang, F. Xu, Global stability of two SIS epidemic mean-field models on complex networks: Lyapunov functional approach, Journal of the Franklin Institute 355 (2018) 6763-6779.
- [461] J. Yang, L. Yang, Z. Jin, Optimal strategies of the age-specific vaccination and antiviral treatment against influenza, Chaos, Solitons & Fractals 168 (2023) 113199.
- [462] L.-X. Yang, X. Yang, J. Liu, Q. Zhu, C. Gan, Epidemics of computer viruses: A complexnetwork approach, Applied Mathematics and Computation 219 (2013) 8705-8717.
- [463] L.-X. Yang, X. Yang, Q. Zhu, L. Wen, A computer virus model with graded cure rates, Nonlinear Analysis: Real World Applications 14 (2013) 414-422.
- [464] L.-X. Yang, X. Yang, The impact of nonlinear infection rate on the spread of computer virus, Nonlinear Dynamics 82 (2015) 85-95.
- [465] Y. Yang, J. Zhou, X. Ma, T. Zhan, Nonstandard finite difference scheme for a diffusive within-host virus dynamics model with both virus-to-cell and cell-to-cell transmissions, Computers & Mathematics with Applications 72 (2016) 1013-1020.
- [466] B. Yu, J. Shi, Z. Xue, M. Yang, X. Yang, Y. Su, Stability analysis of HCV dynamic model with saturation incidence, cellular immunity and interferon effect in intrahepatic and extrahepatic tissues, Mathematics and Computers in Simulation 216 (2024) 301-317.
- [467] Z. U. A. Zafar, M. Inc, F. Tchier, L. Akinyemi, Stochastic suicide substrate reaction model, Physica A: Statistical Mechanics and its Applications 610 (2023) 128384.
- [468] C. Zhang, J. Gao, H. Sun, J. Wang, Dynamics of a reaction-diffusion SVIR model in a spatial heterogeneous environment, Physica A 533 (2019) 122049.

- [469] J. Zhang, J. Lou, Z. Ma, J. Wu, A compartmental model for the analysis of SARS transmission patterns and outbreak control measures in China, Applied Mathematics and Computation 162 (2005) 909-924.
- [470] H. Zhao, L. Wang, S. M. Oliva, H. Zhu, Modeling and Dynamics Analysis of Zika Transmission with Limited Medical Resources, Bulletin of Mathematical Biology (2020) 82:99.
- [471] M. Zhou, H. Xiang, Z. Li, Optimal control strategies for a reaction-diffusion epidemic system, Nonlinear Analysis: Real World Applications 46 (2019) 446-464.
- [472] Q. Zhou, Analysis of opioid transmission crisis model in resilient cities based on numerical solutions of differential equations, Sustainable Energy Technologies and Assessments 52 (2022) 102210.
- [473] C.-C. Zhu, J. Zhu, Dynamic analysis of a delayed COVID-19 epidemic with home quarantine in temporal-spatial heterogeneous via global exponential attractor method, Chaos, Solitons & Fractals 143 (2021) 110546.
- [474] L. Zhu, M. Liu, Y. Li, The dynamics analysis of a rumor propagation model in online social networks, Physica A 520 (2019) 118-137.
- [475] L. Zhu, B. Wang, Stability analysis of a SAIR rumor spreading model with control strategies in online social networks, Information Sciences 526 (2020) 1-19.
- [476] L. Zhu, T. Yuan, Optimal control and parameter identification of a reaction-diffusion network propagation model, Nonlinear Dynamics 111 (2023) 21707-21733.

DEPARTMENT OF MATHEMATICS, FPT UNIVERSITY, HOA LAC HI-TECH PARK, KM29 THANG LONG BLVD, HANOI, VIET NAM

Email address: tuanhm16@fe.edu.vn

UNIVERSITY OF WUPPERTAL, CHAIR OF APPLIED AND COMPUTATIONAL MATHEMATICS, GAUSS-STRASSE 20, 42119 WUPPERTAL, GERMANY

Email address: ehrhardt@uni-wuppertal.de

36